

Supplementary material

Characteristics of included studies

The characteristics of studies included in the review are detailed in tables and appendices within the review. Note that these references are included in the list below along with studies that were rejected upon reading at full text/abstract (reasons for rejection were not recorded).

References retrieved from electronic searching assessed at full text/abstract

- (1993). "Dead wood matters : the ecology and conservation of saproxylic invertebrates in Britain : Meeting : Papers." Peterborough : English Nature.
- (1996). "Thetford Forest Park: the ecology of a pine forest." Forestry Commission Technical Paper 13.
- (1999). "Powis Castle Park Saproxylic invertebrates." CCW contract science report 351.
- (2001). "Welsh parklands survey (stage 5 - 1996) : Llanover Park saproxylic invertebrates." CCW contract science report 354.
- Åkland, B. r., A. Bakke, et al. (1996). "What factors influence the diversity of saproxylic beetles? A multiscaled study from a spruce forest in southern Norway." Biodiversity and Conservation (Historical Archive) 5(1): 75-100.
- Abbott, D. T. and D. A. Crossley, Jr. (1982). "Woody Litter Decomposition Following Clear-Cutting." Ecology 63(1): 35-42.
- Abbott, I., T. Burbidge, et al. (2003). "Logging and burning impacts on cockroaches, crickets and grasshoppers, and spiders in Jarrah forest, Western Australia." Forest Ecology and Management 174(1-3): 383-399.
- Abildsnes, J. and B. A. Tommeras (2000). "Impacts of experimental habitat fragmentation on ground beetles (Coleoptera, Carabidae) in a boreal spruce forest." Annales Zoologici Fennici 37(3): 201-212.
- Acker, S. A., S. V. Gregory, et al. (2003). "Composition, complexity, and tree mortality in riparian forests in the central Western Cascades of Oregon." Forest Ecology and Management 173(1-3): 293-308.
- Addison, J. A., J. A. Trofymow, et al. (2003). "Abundance, species diversity, and community structure of Collembola in successional coastal temperate forests on Vancouver Island, Canada." Applied Soil Ecology 24(3): 233-246.
- Aksoy, A., J. M. Dixon, et al. (1998). "Capsella bursa-pastoris (L.) Medikus (Thlaspi bursa-pastoris L., Bursa bursa-pastoris (L.) Shull, Bursa pastoris (L.) Weber)." Journal of Ecology 86(1): 171-186.
- Alaruikka, D., D. J. Kotze, et al. (2002). "Carabid beetle and spider assemblages along a forested urban-rural gradient in southern Finland." Journal of Insect Conservation 6(4): 195-206.
- Albers, D., S. Migge, et al. (2004). "Decomposition of beech leaves (Fagus sylvatica) and spruce needles (Picea abies) in pure and mixed stands of beech and spruce." Soil Biology and Biochemistry 36(1): 155-164.
- Albrecht, L. (1991). "The Importance of Dead Woody Material in Forests." Forstwissenschaftliches Centralblatt 110(2): 106-113.

- Albrecht, L. (1992). "The Importance of Natural Forest Reserves for Species Protection on Woodlands." Forstwissenschaftliches Centralblatt **111**(4): 214-224.
- Alexander, K. N. A. (1995). "Historic parks and pasture-woodlands: The National Trust resource and its conservation." Biological Journal of the Linnean Society **56**: 155-175.
- Alexander, K. N. A. (1995). "An interesting saproxylic fauna at Snelmore Common, Berkshire." British Journal of Entomology and Natural History **8**(2): 74.
- Alexander, K. N. A. (2002). "Forthampton Oaks, Gloucestershire: a site of major importance for saproxylic invertebrates." British Journal of Entomology and Natural History **15**(1): 63-64.
- Alexander, K. N. A. (2004). "Revision of the Index of ecological continuity as used for saproxylic beetle." English Nature research reports **574**.
- Allen, C. R., R. S. Lutz, et al. (1995). "Red Imported Fire Ant Impacts on Northern Bobwhite Populations." Ecological Applications **5**(3): 632-638.
- Andersen, J., S. Olberg, et al. (2000). "Saproxylic beetles (Coleoptera) of Troms and western Finnmark, northern Norway with exceptional distribution in Fennoscandia." Norwegian Journal of Entomology **47**: 29-40.
- Anderson, J. M. (1991). "The Effects of Climate Change on Decomposition Processes in Grassland and Coniferous Forests." Ecological Applications **1**(3): 326-347.
- Andersson, R. and L. Ostlund (2004). "Spatial patterns, density changes and implications on biodiversity for old trees in the boreal landscape of northern Sweden." Biological Conservation **118**(4): 443-453.
- Andresen, E. (2003). "Effect of forest fragmentation on dung beetle communities and functional consequences for plant regeneration." Ecography **26**(1): 87-97.
- Apolinario, F. E. and C. Martius (2004). "Ecological role of termites (Insecta, Isoptera) in tree trunks in central Amazonian rain forests." Forest Ecology and Management **194**(1-3): 23-28.
- Armbrecht, I., I. Armbrecht, et al. (2003). "Litter-twig dwelling ant species richness and predation potential within a forest fragment and neighboring coffee plantations of contrasting habitat quality in Mexico." Agriculture, Ecosystems and Environment **97**(1-3): 107-115.
- Armstrong, A. J. (2002). "Insects and the determination of priority areas for biodiversity conservation in Kwazulu-Natal province, South Africa." African Entomology **10**(1): 11-27.
- As, S. (1993). "Are Habitat Islands Islands - Woodliving Beetles (Coleoptera) in Deciduous Forest Fragments in Boreal Forest." Ecography **16**(3): 219-228.
- Ashburner, M. and G. Bachli (2004). "Drosophila elegans Bock & Wheeler, 1972 (Insecta, Diptera): proposed conservation of the specific name." Bulletin of Zoological Nomenclature **61**(3): 165-166.
- Ashwini, K. M. and K. R. Sridhar "Leaf litter preference and conversion by a saprophagous tropical pill millipede, *Arthrosphaera magna* Attems." Pedobiologia **In Press, Corrected Proof**.
- Assmann, T. (1999). "The ground beetle fauna of ancient and recent woodlands in the lowlands of north-west Germany (Coleoptera, Carabidae)." Biodiversity and Conservation **8**(11): 1499-1517.

- Assmann, T. and J. Janssen (1999). "The effects of habitat changes on the endangered ground beetle *Carabus nitens* (Coleoptera: Carabidae)." Journal of Insect Conservation **3**(2): 107-116.
- Atilla, N., M. A. Wetzel, et al. (2003). "Abundance and colonization potential of artificial hard substrate-associated meiofauna." Journal of Experimental Marine Biology and Ecology **287**(2): 273-287.
- Atlegrim, O. and K. Sjöberg (1996). "Effects of clear-cutting and single-tree selection harvests on herbivorous insect larvae feeding on bilberry (*Vaccinium myrtillus*) in uneven-aged boreal *Picea abies* forests." Forest Ecology and Management **87**(1-3): 139-148.
- Atlegrim, O. and K. Sjöberg (1996). "Response of bilberry (*Vaccinium myrtillus*) to clear-cutting and single-tree selection harvests in uneven-aged boreal *Picea abies* forests." Forest Ecology and Management **86**(1-3): 39-50.
- Atlegrim, O. and K. Sjöberg (2003). "Patterns of terricolous and field layer invertebrates along a boreal mire-forest gradient." Entomologica Fennica **14**(4): 227-236.
- Atlegrim, O. and K. Sjöberg (2004). "Selective felling as a potential tool for maintaining biodiversity in managed forests." Biodiversity and Conservation **13**(6): 1123-1133.
- Attignon, S. E., D. Weibel, et al. (2004). "Leaf litter breakdown in natural and plantation forests of the Lama forest reserve in Benin." Applied Soil Ecology **27**(2): 109-124.
- Attiwill, P. M. (1994). "The disturbance of forest ecosystems: the ecological basis for conservative management." Forest Ecology and Management **63**(2-3): 247-300.
- Avendañ, C. o-mendoza, et al. Dung beetle community (Coleoptera: Scarabaeidae: Scarabaeinae) in a tropical landscape at the Lachua Region, Guatemala. Biodiversity and Conservation. **14**: 801-822.
- Ayres, M. P. and M. J. Lombardero (2000). "Assessing the consequences of global change for forest disturbance from herbivores and pathogens." The Science of The Total Environment **262**(3): 263-286.
- Baillie, B. R., T. L. Cummins, et al. (1999). "Harvesting effects on woody debris and bank disturbance in stream channels." New Zealand Journal of Forestry Science **29**(1): 85-101.
- Baillie, B. R., T. L. Cummins, et al. (1999). "Measuring woody debris in the small streams of New Zealand's pine plantations." New Zealand Journal of Marine and Freshwater Research **33**(1): 87-97.
- Baillie, B. R. and T. R. Davies (2002). "Influence of large woody debris on channel morphology in native forest and pine plantation streams in the Nelson region, New Zealand." New Zealand Journal of Marine and Freshwater Research **36**(4): 763-774.
- Baker, S. C., A. M. M. Richardson, et al. (2004). "Does clearfell, burn and sow silviculture mimic the effect of wildfire? A field study and review using litter beetles." Forest Ecology and Management **199**(2-3): 433-448.
- Bakke, A. (1999). "High diversity of saproxylic beetles in a hemiboreal mixed forest reserve in the south of Norway." Scandinavian journal of forest research **14**(3): 199-208.
- Bakken, G. S., W. A. Buttemer, et al. (1981). "Heated Taxidermic Mounts: A Means of Measuring the Standard Operative Temperature Affecting Small Animals." Ecology **62**(2): 311-318.

- Baldissera, R., G. Ganade, et al. (2004). "Web spider community response along an edge between pasture and Araucaria forest." Biological Conservation **118**(3): 403-409.
- Ball, I. R., D. B. Lindenmayer, et al. (1999). "A tree hollow dynamics simulation model." Forest Ecology and Management **123**(2-3): 179-194.
- Bamford, M. K., E. M. Roberts, et al. (2002). "An extensive deposit of fossil conifer wood from the Mesozoic of Mali, southern Sahara." Palaeogeography, Palaeoclimatology, Palaeoecology **186**(1-2): 115-126.
- Bancroft, J. S. and P. Turchin (2003). "An experimental test of fragmentation and loss of habitat with *Oryzaephilus surinamensis*." Ecology **84**(7): 1756-1767.
- Bank, R. A. (1999). "On the proposed conservation of *Hydrobia* Hartmann, 1821 (Mollusca, Gastropoda) and *Cyclostoma acutum* Draparnaud, 1805 (currently *Hydrobia acuta*) by the replacement of the lectotype of *H. acuta* with a neotype; proposed designation of *Turbo ventrosus*." Bulletin of Zoological Nomenclature **56**(4): 268-269.
- Barbero, E., C. Palestrini, et al. (1999). "Dung beetle conservation: Effects of habitat and resource selection (Coleoptera: Scarabaeoidea)." Journal of Insect Conservation **3**(2): 75-84.
- Barbosa, O. and P. A. Marquet (2002). "Effects of forest fragmentation on the beetle assemblage at the relict forest of Fray Jorge, Chile." Oecologia **132**(2): 296-306.
- Barkham, J. P. (1978). "Pedunculate Oak Woodland in a Severe Environment: Black Tor Copse, Dartmoor." The Journal of Ecology **66**(3): 707-740.
- Bates, A., J. P. Sadler, et al. (2005). "Spatial dynamics of beetles living on exposed riverine sediments in the upper River Severn: Method development and preliminary results." Aquatic Conservation-Marine and Freshwater Ecosystems **15**(2): 159-174.
- Batzer, D. P. (1998). "Trophic Interactions among Detritus, Benthic Midges, and Predatory Fish in a Freshwater Marsh." Ecology **79**(5): 1688-1698.
- Baur, B., Z. S., et al. Habitat characteristics of the endangered flightless beetle *Dorcadion fuliginator* (Coleoptera: Cerambycidae): implications for conservation. Biological Conservation. **105**: 133-142.
- Bedick, J. C., W. Wyatt Hoback, et al. (1999). "Distribution, ecology, and population dynamics of the American burying beetle [*Nicrophorus americanus* Olivier (coleoptera, silphidae)] in south-central Nebraska, USA." Journal of Insect Conservation **3**(3): 171-181.
- Belausoff, S., P. G. Kevan, et al. (2003). "Assessing tillage disturbance on assemblages of ground beetles (Coleoptera: Carabidae) by using a range of ecological indices." Biodiversity and conservation **12**: 851-882.
- Belloq, M. I. and S. M. Smith (2003). "Convergence in arthropod assemblages with various restoration approaches for Canadian deciduous forests." Journal of Insect Conservation **7**(2): 99-109.
- Bengtsson, J., S. G. Nilsson, et al. (2000). "Biodiversity, disturbances, ecosystem function and management of European forests." Forest Ecology and Management **132**(1): 39-50.
- Benke, A. C., T. C. V. Arsdall, Jr., et al. (1984). "Invertebrate Productivity in a Subtropical Blackwater River: The Importance of Habitat and Life History." Ecological Monographs **54**(1): 25-63.

- Benke, A. C. and D. I. Jacobi (1994). "Production Dynamics and Resource Utilization of Snag-Dwelling Mayflies in a Blackwater River." Ecology **75**(5): 1219-1232.
- Berenzen, N., T. Kumke, et al. (2005). "Macroinvertebrate community structure in agricultural streams: impact of runoff-related pesticide contamination." Ecotoxicology and Environmental Safety **60**(1): 37-46.
- Bergelson, J. M. and J. H. Lawton (1988). "Does Foliage Damage Influence Predation on the Insect Herbivores of Birch?" Ecology **69**(2): 434-445.
- Bergman, K.-O. (1999). "Habitat utilization by Lopinga achine (Nymphalidae: Satyrinae) larvae and ovipositing females: implications for conservation." Biological Conservation **88**(1): 69-74.
- Bess, E. C., R. R. Parmenter, et al. (2002). "Responses of a riparian forest-floor arthropod community to wildfire in the middle Rio Grande Valley, New Mexico." Environmental entomology **31**(5): 774-784.
- Bhattacharya, M. (2004). Restoration ecology and habitat fragmentation studies in Massachusetts woodlands, BOSTON UNIVERSITY.
- Bianchi, F. J. J. A. and W. Van Der Werf (2004). "Model evaluation of the function of prey in non-crop habitats for biological control by ladybeetles in agricultural landscapes." Ecological Modelling **171**(1-2): 177-193.
- Bird, S. B., R. N. Coulson, et al. (2004). "Changes in soil and litter arthropod abundance following tree harvesting and site preparation in a loblolly pine (*Pinus taeda* L.) plantation." Forest Ecology and Management **202**(1-3): 195-208.
- Bishop, D. J. (1998). Saproxyllic beetles and deadwood structure in managed and naturally disturbed spruce forests in Nova Scotia, CARLETON UNIVERSITY (CANADA),.
- Black, H. I. J. and M. J. N. Okwakol (1997). "Agricultural intensification, soil biodiversity and agroecosystem function in the tropics: the role of termites." Applied Soil Ecology **6**(1): 37-53.
- Blake, J. G. (2005). "Effects of prescribed burning on distribution and abundance of birds in a closed-canopy oak-dominated forest, Missouri, USA." Biological Conservation **121**(4): 519-531.
- Blake, S., G. N. Foster, et al. (1994). "Effects of Habitat Type and Grassland Management-Practices on the Body-Size Distribution of Carabid Beetles." Pedobiologia **38**(6): 502-512.
- Blake, S., D. I. McCracken, et al. (2003). "The relationship between the classification of Scottish ground beetle assemblages (Coleoptera, Carabidae) and the National Vegetation Classification of British plant communities." Ecography **26**(5): 602-616.
- Bobiec, A. (2002). "Living stands and dead wood in the Bialowieza forest: suggestions for restoration management." Forest Ecology and Management **165**(1-3): 125-140.
- Bogya, S. and V. Marko (1999). "Effect of pest management systems on ground-dwelling spider assemblages in an apple orchard in Hungary." Agriculture, Ecosystems & Environment **73**(1): 7-18.
- Boling, R. H., Jr., E. D. Goodman, et al. (1975). "Toward A Model of Detritus Processing in a Woodland Stream." Ecology **56**(1): 141-151.
- Bonham, K. J., R. Mesibov, et al. (2002). "Diversity and abundance of some ground-dwelling invertebrates in plantation vs. native forests in Tasmania, Australia." Forest Ecology and Management **158**(1-3): 237-247.

- Bonn, A., K. Hagen, et al. (2002). "The significance of flood regimes for carabid beetle and spider communities in riparian habitats - A comparison of three major rivers in Germany." River Research and Applications **18**(1): 43-64.
- Bonn, A. and M. Kleinwachter (1999). "Microhabitat distribution of spider and ground beetle assemblages (Araneae, Carabidae) on frequently inundated river banks of the River Elbe." Zeitschrift fur Okologie und Naturschutz **8**(3): 109-123.
- Bonn, A. and B. Schroder (2001). "Habitat models and their transfer for single and multi species groups: a case study of carabids in an alluvial forest." Ecography **24**(4): 483-496.
- Boomsma, J. J. and A. J. V. Loon (1982). "Structure and Diversity of Ant Communities in Successive Coastal Dune Valleys." The Journal of Animal Ecology **51**(3): 957-974.
- Boothroyd, I. K. G., J. M. Quinn, et al. (2004). "Riparian buffers mitigate effects of pine plantation logging on New Zealand streams: 1. Riparian vegetation structure, stream geomorphology and periphyton." Forest Ecology and Management **194**(1-3): 199-213.
- Bouget, C. and L. Leseigneur (2005). "The effects of windstorm on saproxylic Eucnemids (Coleoptera, Eucnemidae) in deciduous forests." Bulletin Mensuel-Societe Linneenne De Lyon **74**(3): 81-92.
- Bowles, D. E., C. B. Barr, et al. (2003). "Habitat and phenology of the endangered riffle beetle *Heterelmis comalensis* and a coexisting species, *Microcyloopus pusillus*, (Coleoptera : Elmidae) at Comal Springs, Texas, USA." Archiv Fur Hydrobiologie **156**(3): 361-383.
- Boyle, J. R., J. E. Warila, et al. (1997). "Cumulative effects of forestry practices: An example framework for evaluation from Oregon, U.S.A." Biomass and Bioenergy **13**(4-5): 223-245.
- Braccia, A. and D. P. Batzer (2001). "Invertebrates associated with woody debris in a Southeastern U.S. forested floodplain wetland." Wetlands **21**(1): 18-31.
- Bratton, J. and J. Andrews (1991). "Invertebrate conservation - principles and their application to broad-leaved woodland." British Wildlife **2**(6): 335-344.
- Brouat, C., S. Meusnier, et al. (2004). "Impact of forest management practices on carabids in European fir forests." Forestry **77**(2): 85-97.
- Brown, N., M. Press, et al. (1999). "Growth and Survivorship of Dipterocarp Seedlings: Differences in Shade Persistence Create a Special Case of Dispersal Limitation." Philosophical Transactions: Biological Sciences **354**(1391, Changes and Disturbance in Tropical Rainforest in South-East Asia): 1847-1855.
- Brown, V. K., A. C. Gange, et al. (1987). "The Effect of Insect Herbivory on the Growth and Reproduction of Two Annual *Vicia* Species at Different Stages in Plant Succession." The Journal of Ecology **75**(4): 1173-1189.
- Buchs, W. (2003). "Biodiversity and agri-environmental indicators--general scopes and skills with special reference to the habitat level." Agriculture, Ecosystems & Environment **98**(1-3): 35-78.
- Buckley, G. P., R. Howell, et al. (1997). "Vegetation succession following ride edge management in lowland plantations and woods. 1. The influence of site factors and management practices." Biological Conservation **82**(3): 289-304.

- Buckmann, D. (1959). "Die Auslösung der Umfärbung durch das Hautungshormon bei *Cerura vinula* L. (Lepidoptera, Notodontidae)." Journal of Insect Physiology **3**(2): 159-166.
- Buckner, C. H. and W. J. Turnock (1965). "Avian Predation on the Larch Sawfly, *Pristiphora Erichsonii* (HTG.), (Hymenoptera: Tenthredinidae)." Ecology **46**(3): 223-236.
- Buckton, S. T. and S. J. Ormerod (1997). "Effects of liming on the coleoptera, hemiptera, araneae and opiliones of catchment Wetlands in Wales." Biological Conservation **79**(1): 43-57.
- Buddle, C. M. (2001). "Spiders (Araneae) associated with downed woody material in a deciduous forest in central Alberta, Canada." Agricultural and Forest Entomology **3**(4): 241-251.
- Burgman, M. A. (1996). "Characterisation and delineation of the eucalypt old-growth forest estate in Australia: a review." Forest Ecology and Management **83**(3): 149-161.
- Burke, D. and H. Goulet (1998). "Landscape and area effects on beetle assemblages in Ontario." Ecography **21**(5): 472-479.
- Buse, A. and J. E. G. Good (1993). "The Effects of Conifer Forest Design and Management on Abundance and Diversity of Rove Beetles (Coleoptera, Staphylinidae) - Implications for Conservation." Biological Conservation **64**(1): 67-76.
- Butterfield, J. (1999). "Changes in decomposition rates and Collembola densities during the forestry cycle in conifer plantations." Journal of Applied Ecology **36**(1): 92-100.
- Cambefort, Y. (1985). "Les Coleopteres Scarabaeidae du Parc National de Tai (Cote d'Ivoire)." Revue francaise d'entomologie **7**(5): 337-342.
- Chaney, R. W. (1950). "A Revision of Fossil *Sequoia* and *Taxodium* in Western North America Based on the Recent Discovery of *Metasequoia*." Transactions of the American Philosophical Society **40**(3): 171-263.
- Chapman, M. G. and T. J. Tolhurst (2004). "The relationship between invertebrate assemblages and bio-dependant properties of sediment in urbanized temperate mangrove forests." Journal of Experimental Marine Biology and Ecology **304**(1): 51-73.
- Chen, Z., T. E. Kolb, et al. (2001). "Mechanisms of Douglas-fir resistance to western spruce budworm defoliation: Bud burst phenology, photosynthetic compensation and growth rate." Tree Physiology **21**(16): 1159-1169.
- Christensen, M., K. Hahn, et al. (2005). "Dead wood in European beech (*Fagus sylvatica*) forest reserves." Forest Ecology and Management **210**(1-3): 267-282.
- Chung, A. Y. C., P. Eggleton, et al. (2000). "The diversity of beetle assemblages in different habitat types in Sabah, Malaysia." Bulletin of Entomological Research **90**(6): 475-496.
- Chust, G., J. L. Pretus, et al. (2004). "Scale dependency of insect assemblages in response to landscape pattern." Landscape Ecology **19**(1): 41-57.
- Clark, S. H. E. (2003). "Insect faunas associated with the fossil remains of *Pinus sylvestris* L. in blanket peat from northeast Scotland." Scottish Geographical Journal **119**(1): 39-52.
- Cole, F. R., A. C. Medeiros, et al. (1992). "Effects of the Argentine Ant on Arthropod Fauna of Hawaiian High-Elevation Shrubland." Ecology **73**(4): 1313-1322.

- Cole, L. C. (1946). "A Study of the Cryptozoa of an Illinois Woodland." Ecological Monographs **16**(1): 49-86.
- Cole, L. J., D. I. McCracken, et al. (2002). "Relationships between agricultural management and ecological groups of ground beetles (Coleoptera: Carabidae) on Scottish farmland." Agriculture, Ecosystems and Environment **93**(1-3): 323-336.
- Collier, K. J. and B. R. Baillie (1999). "Decay state and orientation of pinus radiata wood in streams and riparian areas of the central North Island." New Zealand Journal of Forestry Science **29**(2): 225-235.
- Collier, K. J. and J. N. Halliday (2000). "Macroinvertebrate-wood associations during decay of plantation pine in New Zealand pumice-bed streams: Stable habitat or trophic subsidy?" Journal of the North American Benthological Society **19**(1): 94-111.
- Collins, C. S., R. N. Conner, et al. (2002). "Influence of hardwood midstory and pine species on pine bole arthropods." Forest Ecology and Management **164**(1-3): 211-220.
- Cook, S. P. (2002). "Beetle (Coleoptera) diversity in mixed pine-hardwood stands in the Ouachita highlands five years following tree harvests." Southwestern Entomologist **27**(3-4): 269-276.
- Courtney, G. W. (1993). "Archaic black flies and ancient forests: conservation of *Parasimulium* habitats in the Pacific Northwest." Aquatic Conservation **3**(4): 361-373.
- Cowling, J. E., J. I. Spicer, et al. (2003). "Environmental tolerances of an invasive terrestrial amphipod, *Arcitalitrus dorrieni* (Hunt) in Britain." Comparative Biochemistry and Physiology - Part A: Molecular & Integrative Physiology **136**(3): 735-747.
- Craig, M. D. and J. D. Roberts (2005). "The short-term impacts of logging on the jarrah forest avifauna in south-west Western Australia: implications for the design and analysis of logging experiments." Biological Conservation **124**(2): 177-188.
- Creed, I. F., D. L. Morrison, et al. (2004). "Is coarse woody debris a net sink or source of nitrogen in the red spruce - Fraser fir forest of the southern Appalachians, U.S.A.?" Canadian Journal of Forest Research **34**(3): 716-727.
- Creutzburg, V. and G. Mletzko (1969). "Ein Beitrag zur Kaferfauna des Naturschutzgebietes "Ostufer der Muritz" (Sudteil). (Coleoptera: Carabidae, Coccinellidae)." May **16**(1/3): 59-75.
- da Fonseca, G. A. B. and J. G. Robinson (1990). "Forest size and structure: Competitive and predatory effects on small mammal communities." Biological Conservation **53**(4): 265-294.
- Dambach, C. A. (1944). "A Ten-Year Ecological Study of Adjoining Grazed and Ungrazed Woodlands in Northeastern Ohio." Ecological Monographs **14**(3): 255-270.
- Davies, R. G., L. M. Hernández, et al. (2003). "Environmental and spatial influences upon species composition of a termite assemblage across neotropical forest islands." Journal of Tropical Ecology **19**(5): 509-524.
- Davis, A. J. (2000). "Species richness of dung-feeding beetles (Coleoptera: Aphodiidae, Scarabaeidae, Hybosoridae) in tropical rainforest at Danum Valley, Sabah, Malaysia." Coleopterists Bulletin **54**(2): 221-231.

- Davis, A. L. V. (2002). "Dung beetle diversity in South Africa: influential factors, conservation status, data inadequacies and survey design." African Entomology **10**(1): 53-65.
- Davy, A. J., G. F. Bishop, et al. (2001). "Salicornia L. (*Salicornia pusilla* J. Woods, *S. ramosissima* J. Woods, *S. europaea* L., *S. obscura* P.W. Ball & Tutin, *S. nitens* P.W. Ball & Tutin, *S. fragilis* P.W. Ball & Tutin and *S. dolichostachya* Moss)." Journal of Ecology **89**(4): 681-707.
- Day, K. R. and J. Carthy (1988). "Changes in carabid beetle communities accompanying a rotation of sitka spruce." Agriculture, Ecosystems & Environment **24**(4): 407-415.
- de Jong, J., A. Dahlberg, et al. (2004). "Dead wood: A critical resource for maintaining forest biodiversity [Död ved i skogen. Hur mycket behövs för att bevara den biologiska mångfalden?]." Svensk Botanisk Tidskrift **98**(5): 278-297.
- De Warnaffe, G. D. B., M. Dufrière, et al. (2004). "To what extent can management variables explain species assemblages? A study of carabid beetles in forests." Ecography **27**(6): 701-714.
- de Warnaffe, G. D. B. and P. Lebrun (2004). "Effects of forest management on carabid beetles in Belgium: implications for biodiversity conservation." Biological Conservation **118**(2): 219-234.
- De Warnaffe, G. D. B., P. Lebrun, et al. (2004). "Effects of forest management on carabid beetles in Belgium: Implications for biodiversity conservation." Biological Conservation **118**(2): 219-234.
- Deans, A. M., J. R. Malcolm, et al. (2005). "Edge effects and the responses of aerial insect assemblages to structural-retention harvesting in Canadian boreal peatland forests." Forest Ecology and Management **204**(2-3): 249-266.
- Deans, A. M., J. R. Malcolm, et al. (2003). "A comparison of forest structure among old-growth, variable retention harvested, and clearcut peatland black spruce (*Picea mariana*) forests in boreal northeastern Ontario." Forestry Chronicle **79**(3): 579-589.
- DeBruyn, L., S. Thys, et al. (2001). "Effects of vegetation and soil on species diversity of soil dwelling Diptera in a heathland ecosystem." Journal of Insect Conservation **5**(2): 87-97.
- Dejean, A., J. L. Durand, et al. (1996). "Ants inhabiting *Cubitermes* termitaries in African rain forests." Biotropica **28**(4 A): 701-713.
- Delettre, Y. R., N. Morvan, et al. (1998). "Local biodiversity and multi-habitat use in empidoiid flies (Insecta: Diptera, Empidoidea)." Biodiversity and conservation **7**(1): 9-25.
- Dennis, P., D. J., et al. Consequences for biodiversity of reducing inputs to upland temperate pastures: effects on beetles (Coleoptera) of cessation of nitrogen fertilizer application and reductions in stocking rates of sheep. Grass & Forage Science. **59**: 121-135.
- Denton, J. and P. Chandler (2005). "Rotherfield Park, North Hampshire: an important site for saproxylic Coleoptera, Diptera and other insects." British Journal of Entomology and Natural History **18**(1): 9-16.
- Desender, K., A. Erynck, et al. (1999). "Beetle diversity and historical ecology of woodlands in Flanders." Belgian Journal of Zoology **129**(1): 139-155.
- Despots, M., G. Brunet, et al. (2004). "The eastern boreal old-growth balsam fir forest: A distinct ecosystem." Canadian Journal of Botany **82**(6): 830-849.

- Deventer, W. C. V. (1939). "Studies on the Ecology of Secondary Communities in a Deciduous Forest Area." Ecology **20**(2): 198-216.
- Dibble, A. C., J. C. Brissette, et al. (1999). "Putting community data to work: some understory plants indicate red spruce regeneration habitat." Forest Ecology and Management **114**(2-3): 275-291.
- Didham, R. K., P. M. Hammond, et al. (1998). "Beetle Species Responses to Tropical Forest Fragmentation." Ecological Monographs **68**(3): 295-323.
- Didham, R. K., J. H. Lawton, et al. (1998). "Trophic Structure Stability and Extinction Dynamics of Beetles (Coleoptera) in Tropical Forest Fragments." Philosophical Transactions: Biological Sciences **353**(1367): 437-451.
- Dirks-Edmunds, J. C. (1947). "A Comparison of Biotic Communities of the Cedar-Hemlock and Oak-Hickory Associations." Ecological Monographs **17**(3): 235-260.
- Djieto-Lordon, C., A. Dejean, et al. (2004). "Symbiotic mutualism with a community of opportunistic ants: protection, competition, and ant occupancy of the myrmecophyte *Barteria nigritana* (Passifloraceae)." Acta Oecologica **26**(2): 109-116.
- Docherty, M. and S. R. Leather (1997). "Structure and abundance of arachnid communities in Scots and lodgepole pine plantations." Forest Ecology and Management **95**(3): 197-207.
- Donegan, K. K., L. S. Watrud, et al. (2001). "Soil and litter organisms in Pacific northwest forests under different management practices." Applied Soil Ecology **18**(2): 159-175.
- Dowdy, W. W. (1944). "A Community Study of a Disturbed Deciduous Forest Area near Cleveland, Ohio, with Special Reference to Invertebrates." Ecological Monographs **14**(2): 193-222.
- Downie, I. S., J. C. Coulson, et al. (1998). "Distribution of aquatic macroinvertebrates within peatland pool complexes in the Flow Country, Scotland." Hydrobiologia **377**: 95-105.
- Drake, K. J. K. C. M. (1993). "Dead wood matters : the ecology and conservation of saproxylic invertebrates in Britain." English Nature science **7**.
- Driscoll, D. A., D. A. Driscoll, et al. (2005). "Beetle responses to habitat fragmentation depend on ecological traits, habitat condition, and remnant size." Conservation Biology **19**(1): 182-194.
- du Bus de Warnaffe, G. and P. Lebrun (2004). "Effects of forest management on carabid beetles in Belgium: implications for biodiversity conservation." Biological Conservation **118**(2): 219-234.
- Du Plessis, M. (1995). "The effects of fuelwood removal on the diversity of some cavity-using birds and mammals in South Africa." Biological Conservation **74**(2): 77-82.
- Duelli, P. and M. K. Obrist (2003). "Regional biodiversity in an agricultural landscape: the contribution of seminatural habitat islands." Basic and Applied Ecology **4**(2): 129-138.
- Duguay, J. P., P. B. Wood, et al. (2000). "Effects of timber harvests on invertebrate biomass and avian nest success." Wildlife Society Bulletin **28**(4): 1123-1131.
- Dungan, R. J. and D. Kelly (2003). "Effect of host-tree and environmental variables on honeydew production by scale insects (*Ultracoelostoma* sp.) in a high elevation *Nothofagus solandri* forest." New Zealand Journal of Ecology **27**(2): 169-177.

- Dunn, R. R. (2004). "Managing the tropical landscape: a comparison of the effects of logging and forest conversion to agriculture on ants, birds, and lepidoptera." Forest Ecology and Management **191**(1-3): 215-224.
- Dunning, C. E., R. A. Redak, et al. (2003). "Preference and performance of a generalist insect herbivore on *Quercus agrifolia* and *Quercus engelmannii* seedlings from a southern California oak woodland." Forest Ecology and Management **174**(1-3): 593-603.
- Duriez, O., Y. Ferrand, et al. (2005). "Habitat selection of the Eurasian woodcock in winter in relation to earthworms availability." Biological Conservation **122**(3): 479-490.
- Dybas, H. S. and D. D. Davis (1962). "A Population Census of Seventeen-Year Periodical Cicadas (Homoptera: Cicadidae: Magicicada)." Ecology **43**(3): 432-444.
- Dyer, L. E. and D. A. Landis (1997). "Influence of noncrop habitats on the distribution of *Eriborus terebrans* (Hymenoptera: Ichneumonidae) in cornfields." Environmental Entomology **26**(4): 924-932.
- Edgar, W. D. (1971). "The Life-Cycle, Abundance and Seasonal Movement of the Wolf Spider, *Lycosa (Pardosa) lugubris*, in Central Scotland." The Journal of Animal Ecology **40**(2): 303-322.
- Eggleton, P., D. E. Bignell, et al. (2002). "Termite diversity across an anthropogenic disturbance gradient in the humid forest zone of West Africa." Agriculture, Ecosystems & Environment **90**(2): 189-202.
- Eggleton, P., R. Homathevi, et al. (1999). "Termite Assemblages, Forest Disturbance and Greenhouse Gas Fluxes in Sabah, East Malaysia." Philosophical Transactions: Biological Sciences **354**(1391, Changes and Disturbance in Tropical Rainforest in South-East Asia): 1791-1802.
- Ehrlich, P. R. (1996). "Conservation in temperate forests: what do we need to know and do?" Forest Ecology and Management **85**(1-3): 9-19.
- Ellis, W. L. and S. S. Bell (2004). "Canopy gaps formed by mangrove trimming: An experimental test of impact on litter fall and standing litter stock in Southwest Florida (USA)." Journal of Experimental Marine Biology and Ecology **311**(2): 201-222.
- Elton, C. S. (1973). "The Structure of Invertebrate Populations Inside Neotropical Rain Forest." The Journal of Animal Ecology **42**(1): 55-104.
- Elton, C. S. and R. S. Miller (1954). "The Ecological Survey of Animal Communities: With a Practical System of Classifying Habitats by Structural Characters." The Journal of Ecology **42**(2): 460-496.
- Eltz, T., C. A. Brühl, et al. (2003). "Nesting and nest trees of stingless bees (Apidae: Meliponini) in lowland dipterocarp forests in Sabah, Malaysia, with implications for forest management." Forest Ecology and Management **172**(2-3): 301-313.
- Engelmann, F. (1957). "Die Steuerung der Ovarfunktion bei der ovoviviparen Schabe *Leucophaea maderae* (Fabr.)." Journal of Insect Physiology **1**(3): 257-262.
- Engelmark, O., K. Sjöberg, et al. (2001). "Ecological effects and management aspects of an exotic tree species: the case of lodgepole pine in Sweden." Forest Ecology and Management **141**(1-2): 3-13.
- Erwin, T. L. (1995). Arboreal beetles of neotropical forests: Agra Fabricius, the cayennensis complex (Coleoptera: Carabidae: Lebiini: Calleidina).

Carabidology: Population biology and conservation of carabid beetles, Kaunainen; Finland, Finnish Zoological and Botanical Publishing Board.

Escobar, F. (2004). "Diversity and composition of dung beetle (Scarabaeinae) assemblages in a heterogeneous Andean landscape." Tropical Zoology **17**(1): 123-136.

Esseen, P.-A., K.-E. Renhorn, et al. (1996). "Epiphytic Lichen Biomass in Managed and Old-Growth Boreal Forests: Effect of Branch Quality." Ecological Applications **6**(1): 228-238.

Estrada, A. and R. Coates-Estrada (2002). "Dung beetles in continuous forest, forest fragments and in an agricultural mosaic habitat island at Los Tuxtlas, Mexico." Biodiversity and conservation **11**: 1903-1918.

Estrada, A., R. Coates-Estrada, et al. (1998). "Dung and carrion beetles in tropical rain forest fragments and agricultural habitats at Los Tuxtlas, Mexico." Journal of Tropical Ecology **14**(5): 577-593.

Evans, A. M., P. W. Clinton, et al. (2003). "The influence of logs on the spatial distribution of litter-dwelling invertebrates and forest floor processes in New Zealand forests." Forest Ecology and Management **184**(1-3): 251-262.

Eyre, M. D., G. N. Foster, et al. (2003). "An investigation into the relationship between water beetle (Coleoptera) distribution and land cover in Scotland and northeast England." Journal of Biogeography **30**(12): 1835-1849.

Eyre, M. D., D. A. Lott, et al. (1995). Assessing the potential for environmental monitoring using ground beetles (Coleoptera: Carabidae) with riverside and Scottish data. Carabidology: Population biology and conservation of carabid beetles, Kaunainen; Finland, Finnish Zoological and Botanical Publishing Board.

Eyre, M. D., D. A. Lott, et al. (2001). "The Rove Beetles (Coleoptera, Staphylinidae) of Exposed Riverine Sediments in Scotland and Northern England: Habitat Classification and Conservation Aspects." Journal of Insect Conservation **5**(3): 173-186.

Eyre, M. D. and M. L. Luff (1995). "Coleoptera on Post-industrial Land: a Conservation Problem?" Land Contamination and Reclamation **3**(2): 132.

Eyre, M. D. and M. L. Luff (2002). "The use of ground beetles (Coleoptera: Carabidae) in conservation assessments of exposed riverine sediment habitats in Scotland and northern England." Journal of Insect Conservation **6**(1): 25-38.

Eyre, M. D., M. L. Luff, et al. (2001). "The ground beetles (Coleoptera: Carabidae) of exposed riverine sediments in Scotland and northern England." Biodiversity and conservation **10**(3): 403-426.

Eyre, M. D., M. L. Luff, et al. (2003). "The relationship between British ground beetles (Coleoptera, Carabidae) and land cover." Journal of Biogeography **30**(5): 719-730.

Fager, E. W. (1968). "The Community of Invertebrates in Decaying Oak Wood." The Journal of Animal Ecology **37**(1): 121-142.

Fayt, P. (2004). "Old-growth boreal forests, three-toed woodpecker and saproxylic beetles - the importance of landscape management history on local consumer-resource dynamics." Ecological Bulletins- Swedish Natural Science Research Council **51**: 249-258.

Fayt, P., M. M. Machmer, et al. (2005). "Regulation of spruce bark beetles by woodpeckers--a literature review." Forest Ecology and Management **206**(1-3): 1-14.

- Ferguson, S. H. and D. J. Archibald (2002). "The 3/4 power law in forest management: how to grow dead trees." Forest Ecology and Management **169**(3): 283-292.
- Fernandez Fernandez, M. M. and J. M. Salgado Costas (2004). "Recolonization of a burnt pine forest (*Pinus pinaster*) by Carabidae (Coleoptera)." European Journal of Soil Biology **40**(1): 47-53.
- Fichter, E. (1939). "An Ecological Study of Wyoming Spruce-Fir Forest Arthropods with Special Reference to Stratification." Ecological Monographs **9**(2): 183-215.
- Finch, O.-D. (2005). "Evaluation of mature conifer plantations as secondary habitat for epigeic forest arthropods (Coleoptera: Carabidae; Araneae)." Forest Ecology and Management **204**(1): 23-36.
- Fisher, D. O. and I. P. F. Owens (2004). "The comparative method in conservation biology." Trends in Ecology & Evolution **19**(7): 391-398.
- Fleming, T. H. (1985). "Coexistence of Five Sympatric Piper (Piperaceae) Species in a Tropical Dry Forest." Ecology **66**(3): 688-700.
- Ford, H. A., G. W. Barrett, et al. (2001). "Why have birds in the woodlands of Southern Australia declined?" Biological Conservation **97**(1): 71-88.
- Foster, D. R. and G. Motzkin (2003). "Interpreting and conserving the openland habitats of coastal New England: insights from landscape history." Forest Ecology and Management **185**(1-2): 127-150.
- Fournier, E. and M. Loreau (2001). "Respective roles of recent hedges and forest patch remnants in the maintenance of ground-beetle (Coleoptera : Carabidae) diversity in an agricultural landscape." Landscape Ecology **16**(1): 17-32.
- Frampton, G. K., T. Cilgi, et al. (1995). "Effects of grassy banks on the dispersal of some carabid beetles (Coleoptera: Carabidae) on farmland." Biological Conservation **71**(3): 347.
- Francis, J. L., G. F. Peterken, et al. (1996). "Designing, managing and enhancing farm woods to maximise environmental benefits." Aspects of Applied Biology **44**: 1-8.
- Franklin, J. F., T. A. Spies, et al. (2002). "Disturbances and structural development of natural forest ecosystems with silvicultural implications, using Douglas-fir forests as an example." Forest Ecology and Management **155**(1-3): 399-423.
- Freedman, B., V. Zelazny, et al. (1996). "Biodiversity implications of changes in the quantity of dead organic matter in managed forests." Environmental Reviews **4**(3): 238-265.
- Freude, H. (1971). "Gedanken uber Naturschutz und Forstnutzung (Eine Bitte an die zustandigen Forstbehorden um die Erhaltung anbruchiger alter Baume)." **24**(2): 281-287.
- Friauf, J. J. (1953). "An Ecological Study of the Dermaptera and Orthoptera of the Welaka Area in Northern Florida." Ecological Monographs **23**(2): 79-126.
- Fridman, J. and M. Walheim (2000). "Amount, structure, and dynamics of dead wood on managed forestland in Sweden." Forest Ecology and Management **131**(1-3): 23-36.
- Fritz, R. and G. Merriam (1996). "Fencerow and forest edge architecture in eastern Ontario farmland." Agriculture, Ecosystems & Environment **59**(3): 159-170.

- Fritz, R. S. (1983). "Ant Protection of a Host Plant's Defoliator: Consequence of an Ant-Membracid Mutualism." Ecology **64**(4): 789-797.
- Fry, G. and I. Sarlov-Herlin (1997). "The ecological and amenity functions of woodland edges in the agricultural landscape; a basis for design and management." Landscape and Urban Planning **37**(1-2): 45-55.
- Fryar, S. C., I. J. Hodgkiss, et al. (2004). "Succession of fungi on dead and live wood in brackish water in Brunei." Mycologia **96**(2): 219-225.
- Fuller, R. J. and G. F. Peterken (1995). "Woodland and scrub." Managing habitats for conservation: 327-361.
- Futuyma, D. J. and F. Gould (1979). "Associations of Plants and Insects in Deciduous Forest." Ecological Monographs **49**(1): 33-50.
- Gandhi, K. J. K., J. R. Spence, et al. (2001). "Fire residuals as habitat reserves for epigaeic beetles (Coleoptera: Carabidae and Staphylinidae)." Biological Conservation **102**(2): 131-141.
- Gardner, S. M. (1991). "Ground Beetle (Coleoptera, Carabidae) Communities on Upland Heath and Their Association with Heathland Flora." Journal of Biogeography **18**(3): 281-289.
- Gibbons, P., D. B. Lindenmayer, et al. (2002). "Hollow selection by vertebrate fauna in forests of southeastern Australia and implications for forest management." Biological Conservation **103**(1): 1-12.
- Gibbons, P., D. B. Lindenmayer, et al. (1996). "Issues associated with the retention of hollow-bearing trees within eucalypt forests managed for wood production." Forest Ecology and Management **83**(3): 245-279.
- Gibbs, J. P. and E. J. Stanton (2001). "Habitat fragmentation and arthropod community change: Carrion beetles, phoretic mites, and flies." Ecological Applications **11**(1): 79-85.
- Gillespie, R. G. and T. Caraco (1987). "Risk-Sensitive Foraging Strategies of Two Spider Populations." Ecology **68**(4): 887-899.
- Godfray, H. C. J., O. T. Lewis, et al. (1999). "Studying Insect Diversity in the Tropics." Philosophical Transactions: Biological Sciences **354**(1391, Changes and Disturbance in Tropical Rainforest in South-East Asia): 1811-1824.
- Goehring, D. M., G. C. Daily, et al. (2002). "Distribution of ground-dwelling arthropods in tropical countryside habitats." Journal of Insect Conservation **6**(2): 83-91.
- Goldstein, P. Z., P. Z. Goldstein, et al. (2002). "A checklist and commentary on the scarabaeid fauna of the Massachusetts offshore islands (Coleoptera: Scarabaeidae)." Journal of the New York Entomological Society **110**(3-4): 389-401.
- Gonzalez-Ochoa, A. and J. de las Heras (2002). "Effects of post-fire silviculture practices on *Pachyrhinus squamosus* defoliation levels and growth of *Pinus halepensis* Mill." Forest Ecology and Management **167**(1-3): 185-194.
- Götmark, F. and M. Thorell (2003). "Size of nature reserves: Densities of large trees and dead wood indicate high value of small conservation forests in southern Sweden." Biodiversity and Conservation **12**(6): 1271-1285.
- Grant, W. C., Jr. (1956). "An Ecological Study of the Peregrine Earthworm *Pheretima Hupeiensis* in the Eastern United States." Ecology **37**(4): 648-658.
- Gray, M. E. and J. J. Tollefson (1988). "Emergence of the western and northern corn rootworms (Coleoptera: Chrysomelidae) from four tillage systems." Journal of economic entomology **81**(5): 1398-1403.

- Greatorex-davies, J. N., T. H. Sparks, et al. (1994). "The Response of Heteroptera and Coleoptera Species to Shade and Aspect in Rides of Coniferised Lowland Woods in Southern England." Biological Conservation **67**(3): 255-273.
- Green, M., H. D. Mansfield-Williams, et al. (2004). "Reduced hardness as an indicator of susceptibility of timbers to attack by *Euophryum confine* Broun." International Biodeterioration & Biodegradation **53**(1): 33-36.
- Green, P. and G. F. Peterken (1997). "Variation in the amount of dead wood in the woodlands of the Lower Wye Valley, UK in relation to the intensity of management." Forest Ecology and Management **98**(3): 229-238.
- Green, P. W. C. and B. D. Turner (2005). "Food-selection by the booklouse, *Liposcelis bostrychophila* Badonnel (Psocoptera: Liposcelididae)." Journal of Stored Products Research **41**(1): 103-113.
- Greenberg, C. H. and A. McGrane (1996). "A comparison of relative abundance and biomass of ground-dwelling arthropods under different forest management practices." Forest Ecology and Management **89**(1-3): 31-41.
- Grindal, S. D., R. M. Brigham, et al. (1999). "Impacts of forest harvesting on habitat use by foraging insectivorous bats at different spatial scales." Ecoscience **6**(1): 25-34.
- Grove, S. and J. Meggs (2003). "Coarse woody debris, biodiversity and management: A review with particular reference to tasmanian wet eucalypt forests." Australian Forestry **66**(4): 258-272.
- Grove, S. J. (2000). "Trunk window trapping: An effective technique for sampling tropical saproxylic beetles." Memoirs of the Queensland Museum **46**(1): 149-160.
- Grove, S. J. (2001). "Developing appropriate mechanisms for sustaining mature timber habitat in managed natural forest stands." International Forestry Review **3**(4): 272-284.
- Grove, S. J. (2001). "Extent and composition of dead wood in Australian lowland tropical rainforest with different management histories." Forest Ecology and Management **154**(1-2): 35-53.
- Grove, S. J. (2001). "Impacts of forest management on saproxylic beetles in the Australian lowland tropics and the development of appropriate indicators of sustainable forest management." Australian Journal of Entomology **40**(3): 297.
- Grove, S. J. (2002). "The influence of forest management history on the integrity of the saproxylic beetle fauna in an Australian lowland tropical rainforest." Biological Conservation **104**(2): 149-171.
- Grove, S. J. (2002). "Saproxylic insect ecology and the sustainable management of forests." Annual review of ecology and systematics **33**: 1-23.
- Grove, S. J. (2002). "Tree basal area and dead wood as surrogate indicators of saproxylic insect faunal integrity: a case study from the Australian lowland tropics." Ecological Indicators **1**(3): 171-188.
- Grove, S. J. and N. E. Stork (1999). "The conservation of saproxylic insects in tropical forests: A research agenda." Journal of Insect Conservation **3**(2): 67-74.
- Grunner, L. and A. Abud-Antun (1976). "Etude des conditions de sporulation et de conservation d'une souche de *Metarrhizium anisopliae* Sorokin isolée de *Phyllophaga pleei* Bl. en Guadeloupe (Coleoptera: Scarabaeidae)." Turrialba : July/Sept **26**(3): 241-246.

- Gunnarsson, B., K. Nitterus, et al. (2004). "Effects of logging residue removal on ground-active beetles in temperate forests." Forest Ecology and Management **201**(2-3): 229-239.
- Gunther, M. J. and T. R. New (2003). "Exotic Pine Plantations in Victoria, Australia: A Threat to Epigeic Beetle (Coleoptera) Assemblages?" Journal of Insect Conservation **7**(2): 73-84.
- Gutiérrez, D., R. Menéndez, et al. (2004). "Habitat-based conservation priorities for carabid beetles within the Picos de Europa National Park, northern Spain." Biological Conservation **115**(3): 379-393.
- Gutowski, J. M. (1990). "Pollination of the orchid *Dactylorhiza fuchsii* by longhorn beetles in primeval forests of northeastern Poland." Biological conservation **51**(4): 287-297.
- Guyette, R. P. and W. G. Cole (1999). "Age characteristics of coarse woody debris (*Pinus strobus*) in a lake littoral zone." Canadian Journal of Fisheries and Aquatic Sciences **56**(3): 496-505.
- Guyette, R. P. and M. Stambaugh (2003). "The age and density of ancient and modern oak wood in streams and sediments." IAWA Journal **24**(4): 345-353.
- Högl, v. K., et al. Succession of Diptera on dead beech wood: A 10-year study. Pedobiologia. **47**: 61-76.
- Haagvar, S. and B. Oekland (1997). "Saproxylic beetle fauna associated with living sporocarps of *Fomitopsis pinicola* (Fr.) Karst. in four spruce forests with different management histories." Fauna Norvegica Seria B **44**(2): 95-105.
- Halaj, J., R. W. Peck, et al. (2005). "Trophic structure of a macroarthropod litter food web in managed coniferous forest stands: a stable isotope analysis with $[\delta]^{15}N$ and $[\delta]^{13}C$." Pedobiologia **49**(2): 109-118.
- Hall, D. J. (1964). "An Experimental Approach to the Dynamics of a Natural Population of *Daphnia Galeata Mendotae*." Ecology **45**(1): 94-112.
- Hall, J. B. (1992). "Ecology of a key African multipurpose tree species, *Balanites aegyptiaca* (Balanitaceae): the state-of-knowledge." Forest Ecology and Management **50**(1-2): 1-30.
- Hall, R. O., Jr. and J. L. Meyer (1998). "The Trophic Significance of Bacteria in a Detritus-Based Stream Food Web." Ecology **79**(6): 1995-2012.
- Halme, E. and J. Niemela (1993). "Carabid Beetles in Fragments of Coniferous Forest." Annales Zoologici Fennici **30**(1): 17-30.
- Hamer, M. L. and R. H. Slotow (2002). "Conservation application of existing data for South African millipedes (Diplopoda)." African Entomology **10**(1): 29-42.
- Hammond, H. E. J. (1996). Arthropod biodiversity of *Populus* spp. coarse woody material in north-central Alberta, UNIVERSITY OF ALBERTA (CANADA),.
- Hammond, H. E. J., L. D.W., et al. Early colonization of *Populus* wood by saproxylic beetles (Coleoptera). Canadian Journal of Forest Research. **31**: 1175-1183.
- Hammond, H. E. J. and H. E. J. Hammond (1997). "Arthropod biodiversity from *Populus* coarse woody material in north-central Alberta: A review of taxa and collection methods." Canadian Entomologist **129**(6): 1009-1033.
- Hammond, H. E. J., D. W. Langor, et al. (2001). "Early colonization of *Populus* wood by saproxylic beetles (Coleoptera)." Canadian journal of forest research **31**(7): 1175-1183.

- Hammond, H. E. J., D. W. Langor, et al. (2004). "Saproxylic beetles (Coleoptera) using Populus in boreal aspen stands of western Canada: spatiotemporal variation and conservation of assemblages." Canadian journal of forest research **34**: 1-19.
- Hammond, R. B. (1997). "Long-term conservation tillage studies: impact of no-till on seedcorn maggot (Diptera: Anthomyiidae)." Crop protection (Guildford, Surrey) **16**(3): 221-225.
- Hampe, A. and J. Arroyo (2002). "Recruitment and regeneration in populations of an endangered South Iberian Tertiary relict tree." Biological Conservation **107**(3): 263-271.
- Handke, K. and K. Menke (1995). "Ground beetles fauna in reedy areas and fallows: the significance of nature conservation in wet fallow areas in the Bremen river marsh [Laufkaferfauna von Rohrichten und Grünlandbrachen: Naturschutz-Bedeutung feuchter Brachflächen in der Bremer Flussmarsch]." Naturschutz und Landschaftsplanung **27**(3): 106-114.
- Hanel, L. (2001). "Succession of soil nematodes in pine forests on coal-mining sands near Cottbus, Germany." Applied Soil Ecology **16**(1): 23-34.
- Hanel, L. (2003). "Recovery of soil nematode populations from cropping stress by natural secondary succession to meadow land." Applied Soil Ecology **22**(3): 255-270.
- Hanel, L. (2004). "Response of soil nematodes inhabiting spruce forests in the Sumava Mountains to disturbance by bark beetles and clear-cutting." Forest Ecology and Management **202**(1-3): 209-225.
- Hangstromgvar, S. (1999). "Saproxylic beetles visiting living sporocarps of Fomitopsis pinicola and Fomes fomentarius." Norwegian Journal of Entomology **46**: 25-32.
- Hanks, L. M., T. D. Paine, et al. (2005). "Influence of the larval environment on performance and adult body size of the wood-boring beetle Phoracantha semipunctata." Entomologia Experimentalis et Applicata **114**(1): 25-34.
- Hanula, J. L. and D. D. Wade (2003). "Influence of long-term dormant-season burning and fire exclusion on ground-dwelling arthropod populations in longleaf pine flatwoods ecosystems." Forest Ecology and Management **175**(1-3): 163-184.
- Harding, P. (2000). Renewing historic landscapes for the conservation of saproxylic invertebrates in an English parkland. Ecological corridors for invertebrates: strategies of dispersal and recolonisation in today's agricultural and forestry landscapes; Atelier sur les corridors ecologiques pour les invertebres strategies de dispersion et de recolonisation dans le, Neuchatel, Switzerland, Council of Europe.
- Harding, P. T. (1993). Saproxylic Coleoptera as indicators of forest relic biotopes in Lowland Britain. European Invertebrate Survey, Helsinki, WWF.
- Harding, P. T. and K. N. A. Alexander (1992). The saproxylic invertebrates of historic parklands: progress and problems. Dead wood matters: the ecology and conservation of saproxylic invertebrates in Britain, Peterborough.
- Harding, P. T. and K. N. A. Alexander (1994). "The use of saproxylic invertebrates in the selection and evaluation of areas of relic forest in pasture-woodlands." British journal of entomology and natural history **7**(suppl. 1): 21-26.

- HARMON, M. E., J. F. FRANKLIN, et al. (2004). Ecology of Coarse Woody Debris in Temperate Ecosystems
Advances in Ecological Research, Academic Press: 59-234.
- Harmon, M. E., O. N. Krankina, et al. (2000). "Decomposition vectors: A new approach to estimating woody detritus decomposition dynamics." *Canadian Journal of Forest Research* **30**(1): 76-84.
- Harper, K. A., Y. Bergeron, et al. (2005). "Structural development following fire in black spruce boreal forest." *Forest Ecology and Management* **206**(1-3): 293-306.
- Harper, M. J., M. A. McCarthy, et al. (2005). "The abundance of hollow-bearing trees in urban dry sclerophyll forest and the effect of wind on hollow development." *Biological Conservation* **122**(2): 181-192.
- Harris, R. J. and B. R. Burns (2000). "Beetle assemblages of kahikatea forest fragments in a pasture-dominated landscape." *New Zealand Journal of Ecology* **24**(1): 57-67.
- Hartley, M. J. (2002). "Rationale and methods for conserving biodiversity in plantation forests." *Forest Ecology and Management* **155**(1-3): 81-95.
- Hartley, S. E. and T. H. Jones (2003). "Plant diversity and insect herbivores: Effects of environmental change in contrasting model systems." *Oikos* **101**(1): 6-17.
- Hartshorn, G. S. (1995). "Ecological Basis for Sustainable Development in Tropical Forests." *Annual Review of Ecology and Systematics* **26**: 155-175.
- Harz, B. and W. Topp (1999). "Deadwood in commercial forest: A source of danger for the outbreak of pest species?
[Totholz im Wirtschaftswald: Eine gefahrenquelle zur massenvermehrung von schadinsekten?]." *Forstwissenschaftliches Centralblatt* **118**(5): 302-313.
- Hasiotis, S. T. (2003). "Complex ichnofossils of solitary and social soil organisms: understanding their evolution and roles in terrestrial paleoecosystems." *Palaeogeography, Palaeoclimatology, Palaeoecology* **192**(1-4): 259-320.
- Hassell, M. P., J. Latto, et al. (1989). "Seeing the Wood for the Trees: Detecting Density Dependence from Existing Life-Table Studies." *The Journal of Animal Ecology* **58**(3): 883-892.
- Haukioja, E. (2003). "Putting the insect into the birch - Insect interaction." *Oecologia* **136**(2): 161-168.
- Hawkins, C. P., M. L. Murphy, et al. (1982). "Effects of Canopy, Substrate Composition, and Gradient on the Structure of Macroinvertebrate Communities in Cascade Range Streams of Oregon." *Ecology* **63**(6): 1840-1856.
- Hawkins, C. P. and J. R. Sedell (1981). "Longitudinal and Seasonal Changes in Functional Organization of Macroinvertebrate Communities in Four Oregon Streams." *Ecology* **62**(2): 387-397.
- Hedgren, P. O. and L. M. Schroeder (2004). "Reproductive success of the spruce bark beetle *Ips typographus* (L.) and occurrence of associated species: A comparison between standing beetle-killed trees and cut trees." *Forest Ecology and Management* **203**(1-3): 241-250.
- Hedgren, P. O., L. M. Schroeder, et al. (2003). "Tree killing by *Ips typographus* (Coleoptera: Scolytidae) at stand edges with and without colonized felled spruce trees." *Agricultural and Forest Entomology* **5**(1): 67-74.

Heilmann-Clausen, J. and M. Christensen (2004). "Does size matter? On the importance of various dead wood fractions for fungal diversity in Danish beech forests." Forest Ecology and Management **201**(1): 105-117.

Heliola, J., M. Koivula, et al. (2001). "Distribution of carabid beetles (Coleoptera, Carabidae) across a boreal forest-clearcut ecotone." Conservation biology : the journal of the Society for Conservation Biology **15**(2): 370-377.

Hellawell, J. M. (1977). "Change in Natural and Managed Ecosystems: Detection, Measurement and Assessment." Proceedings of the Royal Society of London. Series B, Biological Sciences **197**(1126, A Discussion on the Scientific Aspects of Nature Conservation in Great Britain): 31-56.

Hewitt, N. and M. Kellman (2004). "Factors influencing tree colonization in fragmented forests: an experimental study of introduced seeds and seedlings." Forest Ecology and Management **191**(1-3): 39-59.

Heyborne, W. H., J. C. Miller, et al. (2003). "Ground dwelling beetles and forest vegetation change over a 17-year-period, in western Oregon, USA." Forest Ecology and Management **179**(1-3): 123-134.

Hine, P. M. H. S. J. (1994). "Welsh parklands and their saproxylic invertebrates." CCW contract science report / Countryside Council for Wales **118**.

Hodge, S. J. and G. F. Peterken (1998). "Deadwood in British forests: Priorities and a strategy." Forestry **71**(2): 99-112.

Holdaway, F. G. (1932). "An Experimental Study of the Growth of Populations of the "Flour Beetle" *Tribolium Confusum* Duval, as Affected by Atmospheric Moisture." Ecological Monographs **2**(3): 261-304.

Holloway, A. K. and G. D. Schnell (1997). "Relationship between numbers of the endangered American burying beetle *Nicrophorus americanus* Olivier (Coleoptera: Silphidae) and available food resources." Biological Conservation **81**(1/2): 145-152.

Holmes, P. R., A. P. Fowles, et al. (1993). "The ground beetle (Coleoptera: Carabidae) fauna of Welsh peatland biotopes-species assemblages in relation to peatland habitats and management." Biological Conservation **65**(1): 61.

Hopkins, A. and J. Wainwright (1989). "Changes in botanical composition and agricultural management of enclosed grassland in upland areas of England and Wales, 1970-86, and some conservation implications." Biological Conservation **47**(3): 219-235.

Hopkins, G. W., J. I. Thacker, et al. (2002). "Identifying rarity in insects: The importance of host plant range." Biological Conservation **105**(3): 293-307.

Horne, P. A. and C. L. Edward (1998). "Effects of tillage on pest and beneficial beetles in the Wimmera region of Victoria, Australia." Australian Journal of Entomology **37**(1): 60-63.

Hortal, J., J. M. Lobo, et al. (2001). "Forecasting insect species richness scores in poorly surveyed territories: The case of the Portuguese dung beetles (Col. Scarabaeinae)." Biodiversity and Conservation **10**(8): 1343-1367.

Hou, P. and C. Pan (2001). "Coarse woody debris and its function in forest ecosystem." Chinese Journal of Applied Ecology **12**(2): 309-314.

Hövmeyer, K. and J. Schauer mann (2003). "Succession of Diptera on dead beech wood: A 10-year study." Pedobiologia **47**(1): 61-75.

Hubbard, R. M., J. M. Vose, et al. (2004). "Stand restoration burning in oak-pine forests in the southern Appalachians: Effects on aboveground biomass

and carbon and nitrogen cycling." Forest Ecology and Management **190**(2-3): 311-321.

Hubert, J., V. Jarosik, et al. (2004). "Astigmatid mite growth and fungi preference (Acari: Acaridida): Comparisons in laboratory experiments." Pedobiologia **48**(3): 205-214.

Hughes, J. B., G. C. Daily, et al. (2000). "Conservation of insect diversity: A habitat approach." Conservation Biology **14**(6): 1788-1797.

Hull, H. E., S. Freitag, et al. (1998). "Identification and evaluation of priority conservation areas for Buprestidae (Coleoptera) in South Africa, Lesotho, Swaziland and Namibia." African Entomology **6**(2): 265-274.

Hull, S. D. (2002). The relationships among vegetative structure, arthropod populations, and grassland bird abundance and reproductive success on wildlife production areas in Ohio, THE OHIO STATE UNIVERSITY.

Humphrey, J. W., C. Hawes, et al. (1999). "Relationships between insect diversity and habitat characteristics in plantation forests." Forest Ecology and Management **113**(1): 11-21.

Hunter, M. D. and P. W. Price (1992). "Playing Chutes and Ladders: Heterogeneity and the Relative Roles of Bottom-Up and Top-Down Forces in Natural Communities." Ecology **73**(3): 724-732.

Hurpin, B. and P. H. Robert (1976). "Conservation dans le sol de trois germes pathogenes pour les larves de Melolontha melolontha (Col.: Scarabaeidae)." Entomophaga **21**(1): 73-80.

Hyvarinen, E., J. Kouki, et al. "Short-term effects of controlled burning and green-tree retention on beetle (Coleoptera) assemblages in managed boreal forests." Forest Ecology and Management **In Press, Corrected Proof**.

Ings, T. C. and S. E. Hartley (1999). "The effect of habitat structure on carabid communities during the regeneration of a native Scottish forest." Forest Ecology and Management **119**(1-3): 123-136.

Iperti, G. and I. Hodek (1974). "Induction alimentaire de la dormance imaginaire chez *Semiadalia undecimnotata* Schn. (Coleop. Coccinellidae) pour aider a la conservation des coccinelles elevees au laboratoire avant une utilisation ulterieure." Annales de zoologie: Ecologie animale **6**(1): 41-51.

Irmeler, U., K. Heller, et al. (2002). "Zonation of ground beetles (Coleoptera: Carabidae) and spiders (Araneida) in salt marshes at the North and the Baltic Sea and the impact of the predicted sea level increase." Biodiversity and conservation **11**(7): 1129-1147.

Irmeler, U. and R. Hoernes (2003). "Assignment and evaluation of ground beetle (Coleoptera: Carabidae) assemblages to sites on different scales in a grassland landscape." Biodiversity and conservation **12**: 1405-1419.

Ishii, H. T., S.-I. Tanabe, et al. (2004). "Exploring the relationships among canopy structure, stand productivity, and biodiversity of temperate forest ecosystems." Forest Science **50**(3): 342-355.

Jabin, M., D. Mohr, et al. (2004). "Influence of deadwood on density of soil macro-arthropods in a managed oak-beech forest." Forest Ecology and Management **194**(1): 61-69.

Jaffee, B. A. (2004). "Wood, nematodes, and the nematode-trapping fungus *Arthrobotrys oligospora*." Soil Biology and Biochemistry **36**(7): 1171-1178.

Janzen, D. H. (1973). "Sweep Samples of Tropical Foliage Insects: Effects of Seasons, Vegetation Types, Elevation, Time of Day, and Insularity." Ecology **54**(3): 687-708.

- Jenkins, N. L. and A. A. Hoffmann (2001). "Distribution of *Drosophila serrata* Malloch (Diptera: Drosophilidae) in Australia with particular reference to the southern border." *Australian Journal of Entomology* **40**(1): 41-48.
- Jeyasingh, P. D., C. A. Fuller, et al. (2004). "Habitat-specific life-history variation in the Caribbean termite *Nasutitermes acajutlae* (Isoptera: Termitidae)." *Ecological Entomology* **29**(5): 606-613.
- Johns, A. D. (1984). Ecological effects of selective logging in a west Malaysian rain-forest, Cambridge.
- Johns, A. D. (1985). "Selective logging and wildlife conservation in tropical rain-forest: Problems and recommendations." *Biological Conservation* **31**(4): 355-375.
- Jonas, J. L., M. R. Whiles, et al. (2002). "Aboveground invertebrate responses to land management differences in a central Kansas grassland." *Environmental Entomology* **31**(6): 1142-1152.
- Jones, D. T., P. Eggleton, et al. (2003). "Termite assemblage collapse along a land-use intensification gradient in lowland central Sumatra, Indonesia." *Journal of Applied Ecology* **40**(2): 380-391.
- Jonsell, M. (2004). "Old park trees: A highly desirable resource for both history and beetle diversity." *Journal of Arboriculture* **30**(4): 238-244.
- Jonsell, M., K. Nittérus, et al. (2004). "Saproxylic beetles in natural and man-made deciduous high stumps retained for conservation." *Biological Conservation* **118**(2): 163-173.
- Jonsell, M., G. Nordlander, et al. (1999). "Colonization patterns of insects breeding in wood-decaying fungi." *Journal of Insect Conservation* **3**(2): 145-161.
- Jonsell, M., M. Schroeder, et al. (2003). "The saproxylic beetle *Bolitophagus reticulatus*: Its frequency in managed forests, attraction to volatiles and flight period." *Ecography* **26**(4): 421-428.
- Jonsell, M., M. Schroeder, et al. (2005). "Saproxylic beetles in high stumps of spruce: Fungal flora important for determining the species composition." *Scandinavian journal of forest research* **20**: 54-62.
- Jonsell, M. and J. Weslien (2003). "Felled or standing retained wood - It makes a difference for saproxylic beetles." *Forest Ecology and Management* **175**(1-3): 425-435.
- Jonsell, M., J. Weslien, et al. (1998). "Substrate requirements of red-listed saproxylic invertebrates in Sweden." *Biodiversity and conservation* **7**(6): 749-764.
- Jonsen, I. D. and L. Fahrig (1997). "Response of generalist and specialist insect herbivores to landscape spatial structure." *Landscape Ecology* **12**(3): 185-197.
- Jonsson, B. G. (2000). "Availability of coarse woody debris in a boreal old-growth *Picea abies* forest." *Journal of Vegetation Science* **11**(1): 51-56.
- Jonsson, M., J. Johannesen, et al. (2003). "Comparative genetic structure of the threatened tenebrionid beetle *Oplocephala haemorrhoidalis* and its common relative *Bolitophagus reticulatus*." *Journal of Insect Conservation* **7**(2): 111-124.
- Jonsson, M., O. Kindvall, et al. (2003). "Modelling mating success of saproxylic beetles in relation to search behaviour, population density and substrate abundance." *Animal Behaviour* **65**(6): 1069-1076.

- Jönsson, N., T. Ranius, et al. (2004). "Nutrient richness of wood mould in tree hollows with the Scarabaeid beetle *Osmoderma eremita*." *Animal Biodiversity and Conservation* **27**(2): 79-82.
- Joys, A. C., R. J. Fuller, et al. (2004). "Influences of deer browsing, coppice history, and standard trees on the growth and development of vegetation structure in coppiced woods in lowland England." *Forest Ecology and Management* **202**(1-3): 23-37.
- Judd, S. (1999). "Chirk Castle Park saproxylic invertebrates." *CCW contract science report* **352**.
- Judd, S. (1999). "Powis Castle Park saproxylic invertebrates." *CCW contract science report* **351**.
- Jukes, M. R., R. Ferris, et al. (2002). "The influence of stand structure and composition on diversity of canopy Coleoptera in coniferous plantations in Britain." *Forest Ecology and Management* **163**(1-3): 27-41.
- Jukes, M. R., A. J. Peace, et al. (2001). "Carabid beetle communities associated with coniferous plantations in Britain: the influence of site, ground vegetation and stand structure." *Forest Ecology and Management* **148**(1-3): 271-286.
- Juvonen-Lettington, A. and A. S. Pullin (2001). "The assessment of the saproxylic Coleoptera fauna of lowland parkland and wood pasture: an evaluation of sampling techniques." *Antenna* **25**(2): 97-99.
- Kaila, L., P. Martikainen, et al. (1997). "Dead trees left in clear-cuts benefit saproxylic Coleoptera adapted to natural disturbances in boreal forest." *Biodiversity and conservation* **6**(1): 1-18.
- Kaila, L., P. Martikainen, et al. (1994). "Saproxylic beetles (Coleoptera) on dead birch trunks decayed by different polypore species." *Annales Zoologici Fennici* **31**(1): 97-107.
- Kanowski, J., C. P. Catterall, et al. (2005). "Consequences of broadscale timber plantations for biodiversity in cleared rainforest landscapes of tropical and subtropical Australia." *Forest Ecology and Management* **208**(1-3): 359-372.
- Kanowski, J., C. P. Catterall, et al. (2003). "Development of forest structure on cleared rainforest land in eastern Australia under different styles of reforestation." *Forest Ecology and Management* **183**(1-3): 265-280.
- Kappes, H. and W. Topp (2004). "Emergence of Coleoptera from deadwood in a managed broadleaved forest in central Europe." *Biodiversity and Conservation* **13**(10): 1905-1924.
- Karjalainen, L. and T. Kuuluvainen (2002). "Amount and diversity of coarse woody debris within a boreal forest landscape dominated by *Pinus sylvestris* in Vienansalo wilderness, eastern Fennoscandia." *Silva Fennica* **36**(1): 147-167.
- Kearsley, M. J. C. and T. G. Whitham (1989). "Developmental Changes in Resistance to Herbivory: Implications for Individuals and Populations." *Ecology* **70**(2): 422-434.
- Keddy, P. A. and C. G. Drummond (1996). "Ecological Properties for the Evaluation, Management, and Restoration of Temperate Deciduous Forest Ecosystems." *Ecological Applications* **6**(3): 748-762.
- Keiper, J. B., W. E. Walton, et al. (2002). "Biology and ecology of higher Diptera from freshwater wetlands." *Annual Review of Entomology* **47**: 207-232.

- Kelley, K. C. and K. C. Kelley (1997). "Local diversity of a bioindicator taxon: Tiger beetles (Coleoptera: Cicindelidae: Cicindela spp.) of the big thicket and pineywoods." Texas Journal of Science **49**(3 SUPPL.): 51-66.
- Kelly, J. A. and M. J. Samways (2003). "Diversity and conservation of forest-floor arthropods on a small Seychelles island." Biodiversity and Conservation **12**(9): 1793-1813.
- Kennedy, R. S. H. and T. A. Spies (2005). "Dynamics of hardwood patches in a conifer matrix: 54 years of change in a forested landscape in Coastal Oregon, USA." Biological Conservation **122**(3): 363-374.
- Kevin, G., S. Richard, et al. Urban domestic gardens (II): experimental tests of methods for increasing biodiversity. Biodiversity and Conservation. **14**: 395-413.
- Key, R. (1993). Invertebrate conservation and pollards. Pollard and veteran tree management, Richmond Publishing.
- Key, R. S. (1992). What are saproxylic invertebrates? Dead wood matters: the ecology and conservation of saproxylic invertebrates in Britain, Peterborough.
- Key, R. S. and S. J. Ball (1992). Positive management for saproxylic invertebrates. Dead wood matters: the ecology and conservation of saproxylic invertebrates in Britain, Peterborough.
- Kidd, M. G. and R. W. Longair (1997). "Abundance and diversity of ant (Hymenoptera : Formicidae) assemblages in regenerating forests of northern Saskatchewan." Canadian Field-Naturalist **111**(4): 635-637.
- Kindvall, O. (1999). "Dispersal in a Metapopulation of the Bush Cricket, *Metrioptera bicolor* (Orthoptera: Tettigoniidae)." Journal of Animal Ecology **68**(1): 172-185.
- Kirby, K. J., C. M. Reid, et al. (1998). "Preliminary estimates of fallen dead wood and standing dead trees in managed and unmanaged forests in Britain." Journal of Applied Ecology **35**(1): 148-155.
- Kirby, K. J., R. C. Thomas, et al. (1995). "Pasture-woodland and its conservation in Britain." Biological Journal of the Linnean Society **56**: 135-153.
- Klein, B. C. (1989). "Effects of Forest Fragmentation on Dung and Carrion Beetle Communities in Central Amazonia." Ecology **70**(6): 1715-1725.
- Kleinjan, C. A. and J. K. Scott (1996). "Selection of *Cassida* spp. from southern Africa for the biological control of *Chrysanthemoides monilifera* in Australia." Annals of Applied Biology **128**(3): 373-385.
- Kless, J. (1974). "Die Kaferarten des Schutzgebietes "Taubergiessen" am Oberrhein." **7**: 552-569.
- Kluwer (2001). "Biodiversity and conservation of diptera." Selected papers. Held at the 4th international congress of dipterology.
- Knight, D. H., W. L. Baker, et al. (2001). "A landscape perspective on the establishment of exotic tree plantations: lodgepole pine (*Pinus contorta*) in Sweden." Forest Ecology and Management **141**(1-2): 131-142.
- Koch, S. O., A. S. Van Jaarsveld, et al. (2000). "Conservation strategies for poorly surveyed taxa: A dung beetle (Coleoptera, Scarabaeidae) case study from southern Africa." Journal of Insect Conservation **4**(1): 45-56.
- Kohli, L., O. Daniel, et al. (1999). "Miscanthus sinensis and wild flowers as food resources of *Lumbricus terrestris* L." Applied Soil Ecology **11**(2-3): 189-197.

- Koivula, M. (2002). "Alternative harvesting methods and boreal carabid beetles (Coleoptera, Carabidae)." Forest Ecology and Management **167**(1-3): 103-121.
- Koivula, M., V. Hyyryläinen, et al. (2004). "Carabid beetles (Coleoptera: Carabidae) at forest-farmland edges in southern Finland." Journal of Insect Conservation **8**(4): 297-309.
- Koivula, M., J. Kukkonen, et al. (2002). "Boreal carabid-beetle (Coleoptera, Carabidae) assemblages along the clear-cut originated succession gradient." Biodiversity and conservation **11**(7): 1269-1288.
- Koivula, M. and J. Niemela (2002). "Boreal carabid beetles (Coleoptera, Carabidae) in managed spruce forests - a summary of Finnish case studies." Silva Fennica **36**(1): 423-436.
- Kolstrom, M. and J. Lumatjarvi (1999). "Decision support system for studying effect of forest management on species richness in boreal forests." Ecological Modelling **119**(1): 43-55.
- Kolstrom, M. and J. Lumatjarvi (2000). "Saproxylic beetles on aspen in commercial forests: a simulation approach to species richness." Forest ecology and management **126**(2): 113-120.
- Komonen, A. (2001). "Structure of insect communities inhabiting old-growth forest specialist bracket fungi." Ecological Entomology **26**(1): 63-75.
- Komonen, A. (2003). "Hotspots of Insect Diversity in Boreal Forests." Conservation Biology **17**(4): 976-981.
- Kotze, D. J., J. Niemela, et al. (2000). "Colonization success of carabid beetles on Baltic islands." Journal of Biogeography **27**(4): 807-819.
- Kotze, D. J., J. Niemela, et al. (2003). "Testing abundance-range size relationships in European carabid beetles (Coleoptera, Carabidae)." Ecography **26**(5): 553-566.
- Kotze, D. J. and R. B. O'Hara (2003). "Species decline - But why? Explanations of carabid beetle (Coleoptera, Carabidae) declines in Europe." Oecologia **135**(1): 138-148.
- Kotze, D. J. and M. J. Samways (2001). "No general edge effects for invertebrates at Afriomontane forest/grassland ecotones." Biodiversity and Conservation **10**(3): 443-466.
- Kouki, J., K. Arnold, et al. (2004). "Long-term persistence of aspen - A key host for many threatened species - Is endangered in old-growth conservation areas in Finland." Journal for Nature Conservation **12**(1): 41-52.
- Kouki, J., S. Lofman, et al. (2001). "Forest fragmentation in Fennoscandia: Linking habitat requirements of wood-associated threatened species to landscape and habitat changes." Scandinavian Journal of Forest Research: 27-37.
- Kramm, K. R., D. F. West, et al. (1982). "Termite pathogens: Transfer of the entomopathogen *Metarhizium anisopliae* between *Reticulitermes* sp. termites." Journal of Invertebrate Pathology **40**(1): 1-6.
- Krankina, O. N., M. E. Harmon, et al. (1999). "Nutrient stores and dynamics of woody detritus in a boreal forest: Modeling potential implications at the stand level." Canadian Journal of Forest Research **29**(1): 20-32.
- Kratochwil, A. (1983). "Zur Phanologie von Pflanzen und blütenbesuchenden Insekten (Hymenoptera, Lepidoptera, Diptera, Coleoptera) eines versauerten Halbtrockenrasens im Kaiserstuhl--ein Beitrag zur Erhaltung brachliegender wiesen als Lizenz-Biotope gefährdeter Tierarten." Veröffentlichungen für

- Naturschutz und Landschaftspflege in Baden Wurttemberg : Beiheft 34: 57-108.
- Kremen, C. (1992). "Assessing the Indicator Properties of Species Assemblages for Natural Areas Monitoring." Ecological Applications **2**(2): 203-217.
- Kromp, B. (1999). "Carabid beetles in sustainable agriculture: a review on pest control efficacy, cultivation impacts and enhancement." Agriculture, Ecosystems & Environment **74**(1-3): 187-228.
- Kubes, J. (1996). "Biocentres and corridors in a cultural landscape. A critical assessment of the 'territorial system of ecological stability'." Landscape and Urban Planning **35**(4): 231-240.
- Kurttila, M., J. Uuttera, et al. (2002). "Decreasing the fragmentation of old forests in landscapes involving multiple ownership in Finland: economic, social and ecological consequences." Forest Ecology and Management **166**(1-3): 69-84.
- Kuschel, G. (1960). "Terrestrial Zoology in Southern Chile." Proceedings of the Royal Society of London. Series B, Biological Sciences **152**(949, A Discussion on the Biology of the Southern Cold Temperate Zone): 540-550.
- Kuuluvainen, T. (2002). "Natural variability of forests as a reference for restoring and managing biological diversity in boreal Fennoscandia." Silva Fennica **36**(1): 97-125.
- Laaka, S. (1992). "The threatened epixylic bryophytes in old primeval forests in Finland." Biological Conservation **59**(2-3): 151-154.
- Laiolo, P., E. Caprio, et al. (2003). "Effects of logging and non-native tree proliferation on the birds overwintering in the upland forests of north-western Italy." Forest Ecology and Management **179**(1-3): 441-454.
- Lambert, F. R. (1992). "The Consequences of Selective Logging for Bornean Lowland Forest Birds." Philosophical Transactions: Biological Sciences **335**(1275, Tropical Rain Forest: Disturbance and Recovery): 443-457.
- Landis, D. A., S. D. Wratten, et al. (2000). "Habitat management to conserve natural enemies of arthropod pests in agriculture." Annual Review of Entomology **45**: 175-201.
- Landres, P. B. and J. A. MacMahon (1983). "Community Organization of Arboreal Birds in Some Oak Woodlands of Western North America." Ecological Monographs **53**(2): 183-208.
- Lane, M. B. and G. McDonald (2002). "Crisis, change, and institutions in forest management: the Wet Tropics of northeastern Australia." Journal of Rural Studies **18**(3): 245-256.
- Lang, G. E. and R. T. T. Forman (1978). "Detrital Dynamics in a Mature Oak Forest: Hutcheson Memorial Forest, New Jersey." Ecology **59**(3): 580-595.
- Langor, D. W., S. J.R., et al. Saproxylic beetles (Coleoptera) using Populus in boreal aspen stands of western Canada: spatiotemporal variation and conservation of assemblages. Canadian Journal of Forest Research. **34**: 1-19.
- Larsen, K. J. and T. W. Work (2003). "Differences in ground beetles (Coleoptera: Carabidae) of original and reconstructed tallgrass prairies in northeastern Iowa, USA, and impact of 3-year spring burn cycles." Journal of Insect Conservation **7**(3): 153-166.
- Larson, M. A., F. R. Thompson III, et al. (2004). "Linking population viability, habitat suitability, and landscape simulation models for conservation planning." Ecological Modelling **180**(1): 103-118.

- Lassau, S. A., D. F. Hochuli, et al. (2005). "Effects of habitat complexity on forest beetle diversity: Do functional groups respond consistently?" Diversity and Distributions **11**(1): 73-82.
- Laurance, W. F., T. E. Lovejoy, et al. (2002). "Ecosystem decay of Amazonian forest fragments: A 22-year investigation." Conservation Biology **16**(3): 605-618.
- Lazdinis, M. and P. Angelstam (2005). "Functionality of riparian forest ecotones in the context of former Soviet Union and Swedish forest management histories." Forest Policy and Economics **7**(3): 321-332.
- Lee, P. C., S. Crites, et al. (1997). "Characteristics and Origins of Deadwood Material in Aspen-Dominated Boreal Forests." Ecological Applications **7**(2): 691-701.
- Lee, P. C. and M. D. Hauser (1998). "Long-Term Consequences of Changes in Territory Quality on Feeding and Reproductive Strategies of Vervet Monkeys." Journal of Animal Ecology **67**(3): 347-358.
- Lefko, S. A., L. P. Pedigo, et al. (1998). "Spatial modeling of preferred wireworm (Coleoptera: Elateridae) habitat." Environmental entomology **27**(2): 184-190.
- Lefko, S. A., L. P. Pedigo, et al. (1998). "Wireworm (Coleoptera: Elateridae) incidence and diversity in Iowa conservation reserve environments." Environmental entomology **27**(2): 312-317.
- Lemieux, J. P. and B. S. Lindgren (2004). "Ground beetle responses to patch retention harvesting in high elevation forests of British Columbia." Ecography **27**(5): 557-566.
- Lima, A. P., A. C. Cordeiro-Duarte, et al. (2000). "Effect of selective logging intensity on two termite species of the genus *Syntermes* in Central Amazonia." Forest Ecology and Management **137**(1-3): 151-154.
- Lindbladh, M., M. Niklasson, et al. (2003). "Long-time record of fire and open canopy in a high biodiversity forest in southeast Sweden." Biological Conservation **114**(2): 231-243.
- Lindenmayer, D. B., R. B. Cunningham, et al. (1997). "Decay and Collapse of Trees with Hollows in Eastern Australian Forests: Impacts on Arboreal Marsupials." Ecological Applications **7**(2): 625-641.
- Lindenmayer, D. B., R. B. Cunningham, et al. (2000). "Structural features of old-growth Australian montane ash forests." Forest Ecology and Management **134**(1-3): 189-204.
- Lindenmayer, D. B., R. D. Incoll, et al. (1999). "Attributes of logs on the floor of Australian Mountain Ash (*Eucalyptus regnans*) forests of different ages." Forest Ecology and Management **123**(2-3): 195-203.
- Lindenmayer, D. B. and H. F. Recher (1998). "Aspects of ecologically sustainable forestry in temperate eucalypt forests - beyond an expanded reserve system." Pacific Conservation Biology **4**(1): 4-10.
- Linder, P., O. Zackrisson, et al. (1997). "Stand structure and successional trends in virgin boreal forest reserves in Sweden." Forest Ecology and Management **98**(1): 17-33.
- Lindgren, M. (2001). "Polypore (Basidiomycetes) species richness and community structure in natural boreal forests of NW Russian Karelia and adjacent areas in Finland." Acta Botanica Fennica(170): 1-41.

- Lindhe, A. (2004). Conservation through management: Cut wood as substrate for saproxylic organisms, SVERIGES LANTBRUKSUNIVERSITET (SWEDEN),.
- Lindhe, A., N. A?senblad, et al. (2004). "Cut logs and high stumps of spruce, birch, aspen and oak - Nine years of saproxylic fungi succession." Biological Conservation **119**(4): 443-454.
- Lindhe, A. and A. Lindelow (2004). "Cut high stumps of spruce, birch, aspen and oak as breeding substrates for saproxylic beetles." Forest ecology and management **13**(203): 1-3.
- Lobo, J. M. (2001). "Decline of roller dung beetle (Scarabaeinae) populations in the Iberian peninsula during the 20th century." Biological Conservation **97**(1): 43-50.
- Lobo, J. M., J.-P. Lumaret, et al. (1997). "Les atlas faunistiques comme outils d'analyse spatiale de la biodiversité." Annales de la Societe Entomologique de France **33**(2): 129-138.
- Londo, M., J. Dekker, et al. (2005). "Willow short-rotation coppice for energy and breeding birds: an exploration of potentials in relation to management." Biomass and Bioenergy **28**(3): 281-293.
- Lovei, G. L. and M. Cartellieri (2000). "Ground beetles (Coleoptera, Carabidae) in forest fragments of the Manawatu, New Zealand: Collapsed assemblages?" Journal of Insect Conservation **4**(4): 239-244.
- Lucas, S. (1988). "Spiders in Brazil." Toxicon **26**(9): 759-772.
- Luscher, M. and P. Karlson (1958). "Experimentelle auslosung von Hautungen bei der Termite *Kaloterme flavicollis* (Fabr.)." Journal of Insect Physiology **1**(4): 341-345.
- Mac Nally, R., A. Parkinson, et al. (2001). "Relationships between terrestrial vertebrate diversity, abundance and availability of coarse woody debris on south-eastern Australian floodplains." Biological Conservation **99**(2): 191-205.
- Mac Nally, R., T. R. Soderquist, et al. (2000). "The conservation value of mesic gullies in dry forest landscapes: Avian assemblages in the box-ironbark ecosystem of southern Australia." Biological Conservation **93**(3): 293-302.
- MacGowan, I. "The entomological value of aspen in the Scottish Highlands." Malloch Society research report **1**.
- Mackay, R. J. and J. Kalff (1969). "Seasonal Variation in Standing Crop and Species Diversity of Insect Communities in a small Quebec Stream." Ecology **50**(1): 101-109.
- MacNally, R., A. Parkinson, et al. (2002). "Current loads of coarse woody debris on southeastern Australian floodplains: Evaluation of change and implications for restoration." Restoration Ecology **10**(4): 627-635.
- Maelfait, J. P., G. Coulon, et al. (1991). The Flemish legislation on the protection of Coleoptera in the European perspective. Faunal inventories of sites for cartography and nature conservation, Brussels, Brussels.
- Maeto, K., S. Sato, et al. (2002). "Species diversity of longicorn beetles in humid warm-temperate forests: the impact of forest management practices on old-growth forest species in southwestern Japan." Biodiversity and conservation **11**: 1919-1937.
- Magura, T., Z. Bordán, et al. (2000). "Effects of nature management practice on carabid assemblages (Coleoptera: Carabidae) in a non-native plantation." Biological Conservation **93**(1): 95-102.

- Magura, T., V. Ködöböcz, et al. (2001). "Effects of forestry practices on carabids (Coleoptera: Carabidae) - implications for nature management." Acta Phytopathologica et Entomologica Hungarica **36**(1-2): 179-188.
- Magura, T., V. Kodobocz, et al. (2001). "Effects of habitat fragmentation on carabids in forest patches." Journal of Biogeography **28**(1): 129-138.
- Magura, T., B. Tóthmezei, et al. (2001). "Forest edge and diversity: Carabids along forest-grassland transects." Biodiversity and Conservation **10**(2): 287-300.
- Maheshwari, G. and H. R. Sou (2000). The Importance of Diptera (Insecta) in Conservation of the Nature with Special Reference to Northwest Himalaya, India. National symposium on environmental protection, Jodhpur, India, [publ date], Jaipur.
- Majer, J. D., H. F. Recher, et al. (1994). "Comparison of arthropod species richness in eastern and western Australian canopies: a contribution to the species number debate." Memoirs - Queensland Museum **36**(1): 121-131.
- Major, R. E., F. J. Christie, et al. (2003). "The effect of habitat configuration on arboreal insects in fragmented woodlands of south-eastern Australia." Biological Conservation **113**(1): 35-48.
- Major, R. E., D. Smith, et al. (1999). "Are roadside strips important reservoirs of invertebrate diversity? A comparison of the ant and beetle faunas of roadside strips and large remnant woodlands." Australian Journal of Zoology **47**(6): 611-624.
- Mallory, E. C., A. M. Gordon, et al. (2000). "Distribution of woody debris in a small headwater lake, central Ontario, Canada." Archiv fur Hydrobiologie **148**(4): 587-606.
- Manna, M. C., S. Jha, et al. (2003). "Comparative efficacy of three epigeic earthworms under different deciduous forest litters decomposition." Bioresource Technology **88**(3): 197-206.
- Marinho da Costa, R. and F. Fernandez (2002). "Feeding and survival rates of the copepods *Euterpina acutifrons* Dana and *Acartia grani* Sars on the dinoflagellates *Alexandrium minutum* Balech and *Gyrodinium corsicum* Paulmier and the Chytridophyta *Rhodomonas baltica* Karsten." Journal of Experimental Marine Biology and Ecology **273**(2): 131-142.
- Marino, P. C. and D. A. Landis (1996). "Effect of Landscape Structure on Parasitoid Diversity and Parasitism in Agroecosystems." Ecological Applications **6**(1): 276-284.
- Marinoni Renato, C. G. N. G. (2003). "Sazonalidade de *Nyssodrysinia lignaria* (Bates) (Coleoptera, Cerambycidae, Lamiinae), no Estado do Paraná, Brasil." Revista Brasileira de Zoologia **20**(1): 141-152.
- Marshall, E. J. P. and A. C. Moonen (2002). "Field margins in northern Europe: their functions and interactions with agriculture." Agriculture, Ecosystems & Environment **89**(1-2): 5-21.
- Marshall, V. G. (2000). "Impacts of forest harvesting on biological processes in northern forest soils." Forest Ecology and Management **133**(1-2): 43-60.
- Martikainen, P. (1999). Conservation of threatened saproxylic beetles: significance of retained aspen *Populus tremula* on clearcut areas. Ecology of coarse woody debris in boreal forests, Oxford.
- Martikainen, P. (2001). "Conservation of threatened saproxylic beetles: significance of retained aspen *Populus tremula* on clearcut areas." Ecological Bulletins- Swedish Natural Science Research Council **49**: 205-218.

- Martikainen, P. (2002). "Ecology and conservation status of *Acanthocinus griseus* (Fabricius, 1792) (Coleoptera: Cerambycidae) in Finland." Entomologica Fennica **13**(1): 41-50.
- Martikainen, P. and L. Kaila (2004). "Sampling saproxylic beetles: Lessons from a 10-year monitoring study." Biological Conservation **120**(2): 175-185.
- Martikainen, P., L. Kaila, et al. (1998). "Threatened beetles in White-backed Woodpecker habitats." Conservation biology : the journal of the Society for Conservation Biology **12**(2): 293-301.
- Martikainen, P. and J. Kouki (2003). "Sampling the rarest: threatened beetles in boreal forest biodiversity inventories." Biodiversity and conservation **12**: 1815-1831.
- Martikainen, P., J. Siitonen, et al. (1999). "Bark beetles (Coleoptera, Scolytidae) and associated beetle species in mature managed and old-growth boreal forests in southern Finland." Forest Ecology and Management **116**(1-3): 233-245.
- Martikainen, P., J. Siitonen, et al. (2000). "Species richness of Coleoptera in mature managed and old-growth boreal forests in southern Finland." Biological Conservation **94**(2): 199-209.
- Martin, D. and M. Uhlig (1986). "Die Spinnen- und Kurzfluglerfauna (Araneae et Staphylinidae) der Silbergrasrasen (Corynephoreten) des Bühnenwerder, Stadtkreis Brandenburg, Bezirk Potsdam (Arachnida; Insecta, Coleoptera)." Faunistische Abhandlungen **14**: 31-35.
- Martin, K. and J. M. Eadie (1999). "Nest webs: A community-wide approach to the management and conservation of cavity-nesting forest birds." Forest Ecology and Management **115**(2-3): 243-257.
- Mason, C. F., S. M. Macdonald, et al. (1984). "Structure, management and conservation value of the riparian woody plant community." Biological Conservation **29**(3): 201-216.
- Mathews, C. R., D. G. Bottrell, et al. (2002). "A comparison of conventional and alternative understory management practices for apple production: multi-trophic effects." Applied Soil Ecology **21**(3): 221-231.
- Maudsley, M. J. (2000). "A review of the ecology and conservation of hedgerow invertebrates in Britain." Journal of Environmental Management **60**(1): 65-76.
- Mawdsley, J. R. (1890). "Ecology, biogeography, and conservation of checkered beetles (Insecta: Coleoptera: Cleridae) in southeastern Arizona: a geographic information system (GIS) study." Transactions of the American Entomological Society: 3.
- McCarthy, M. A. and D. B. Lindenmayer (1998). "Multi-aged mountain ash forest, wildlife conservation and timber harvesting." Forest Ecology and Management **104**(1-3): 43-56.
- McCarty, J. P., D. J. Levey, et al. (2002). "Spatial and temporal variation in fruit use by wildlife in a forested landscape." Forest Ecology and Management **164**(1-3): 277-291.
- McCay, T. S. and M. J. Komoroski (2004). "Demographic responses of shrews to removal of coarse woody debris in a managed pine forest." Forest Ecology and Management **189**(1-3): 387-395.
- McClure, J. M., R. K. Kolka, et al. (2004). "Effect of forest harvesting best management practices on coarse woody debris distribution in stream and

riparian zones in three Appalachian watersheds." Water, Air, and Soil Pollution: Focus **4**(1): 245-261.

McCracken, D. I., I. S. Downie, et al. (2005). "Comparing the effects of farming practices on ground beetle (Coleoptera: Carabidae) and spider (Araneae) assemblages of Scottish farmland." Biodiversity and Conservation **14**(2): 441-460.

McGee, G. G., D. J. Leopold, et al. (1999). "Structural Characteristics of Old-Growth, Maturing, and Partially Cut Northern Hardwood Forests." Ecological Applications **9**(4): 1316-1329.

McHugh, J. V. (1995). "Aspidiphorus Ziegler in Dejean, 1821 (Insecta, Coleoptera): proposed conservation as the correct original spelling, and Aspidiphoridae Kiesenwetter, 1877 (1859): proposed placement on the Official List." Bulletin of Zoological Nomenclature **52**(1): 44.

McIntyre, N. E. and T. R. Thompson (2003). "A Comparison of Conservation Reserve Program Habitat Plantings with Respect to Arthropod Prey for Grassland Birds." American Midland Naturalist **150**(2): 291-301.

McIver, J. D., T. Torgersen, et al. (2003). "Environmental Effects and Economics of Mechanized Logging for Fuel Reduction in Northeastern Oregon Mixed-Conifer Stands." Western Journal of Applied Forestry **18**(4): 238-249.

McLean, I. F. G. and M. C. D. Speight (1992). Saproxylic invertebrates - the European context. Dead wood matters: the ecology and conservation of saproxylic invertebrates in Britain, Peterborough.

McWilliam, H. A., R. G. Death, et al. (1998). "Arboreal arthropod communities of remnant podocarp-hardwood rainforest in North Island, New Zealand." New Zealand Journal of Zoology **25**(2): 157-169.

Meggs, J. M., S. A. Munks, et al. (2004). "Development and evaluation of predictive habitat models to assist the conservation planning of a threatened lucanid beetle, *Hoplogonus simsoni*, in north-east Tasmania." Biological Conservation **118**(4): 501-511.

Meggs, J. M., S. A. Munks, et al. (2003). "Distribution, habitat characteristics and conservation requirements of a forest-dependent threatened invertebrate *Lissotes latidens* (Coleoptera: Lucanidae)." Journal of Insect Conservation **7**(3): 137-152.

Meggs, J. M., S. A. Munks, et al. (2003). "The distribution and habitat characteristics of a threatened lucanid beetle *Hoplogonus simsoni* in north-east Tasmania." Pacific Conservation Biology **9**(3): 172-186.

Meggs, J. M. and R. J. Taylor (1999). "Distribution and conservation status of the Mt Mangana stag beetle, *Lissotes menalcas* (Coleoptera: Lucanidae)." Papers and Proceedings- Royal Society of Tasmania **133**(1): 23-28.

Meleason, M. A. and J. M. Quinn (2004). "Influence of riparian buffer width on air temperature at Whangapoua Forest, Coromandel Peninsula, New Zealand." Forest Ecology and Management **191**(1-3): 365-371.

Menno, R. Saproxylic hoverflies benefit by modern forest management (Diptera: Syrphidae). Journal of Insect Conservation. **9**: 49-59.

Mensah, R. K. and J. L. Madden (1994). "Conservation of two predator species for biological control of *Chrysophtharta bimaculata* (Col.: Chrysomelidae) in Tasmanian forests." Entomophaga **39**(1): 71-83.

- Meyer, S. G. E. and G. Schaub (1973). "Der respiratorische Stoffwechsel von Calliphoridenlarven in Beziehung zu Temperaturadaptation und Regulation." Journal of Insect Physiology **19**(11): 2183-2187.
- Michaels, K. and G. Bornemissza (1999). "Effects of clearfell harvesting on lucanid beetles (Coleoptera: Lucanidae) in wet and dry sclerophyll forests in Tasmania." Journal of Insect Conservation **3**(2): 85-95.
- Middleton, J. and G. Merriam (1985). "The rationale for conservation: Problems from a virgin forest." Biological Conservation **33**(2): 133-145.
- Minarro, M. and E. Dapena (2003). "Effects of groundcover management on ground beetles (Coleoptera: Carabidae) in an apple orchard." Applied Soil Ecology **23**(2): 111-117.
- Monserud, R. A. (2002). "Large-scale management experiments in the moist maritime forests of the Pacific Northwest." Landscape and Urban Planning **59**(3): 159-180.
- Moore, J. D., R. Ouimet, et al. (2004). "Effects of two silvicultural practices on ground beetles (Coleoptera : Carabidae) in a northern hardwood forest, Quebec, Canada." Canadian Journal of Forest Research-Revue Canadienne De Recherche Forestiere **34**(4): 959-968.
- Moore, J.-D., R. Ouimet, et al. (2002). "Effects of two silvicultural practices on soil fauna abundance in a northern hardwood forest, Québec, Canada." Canadian Journal of Soil Science **82**(1): 105-113.
- Moore, J. L., S. M. Howden, et al. (2001). "The dynamics of grazed woodlands in southwest Queensland, Australia, and their effect on greenhouse gas emissions." Environment International **27**(2-3): 147-153.
- Moran, V. C. and T. R. E. Southwood (1982). "The Guild Composition of Arthropod Communities in Trees." The Journal of Animal Ecology **51**(1): 289-306.
- Moretti, M. and S. Barbalat (2004). "The effects of wildfires on wood-eating beetles in deciduous forests on the southern slope of the Swiss Alps." Forest Ecology and Management **187**(1): 85-103.
- Morgan, M., C. B. Knisley, et al. (2000). "New taxonomic status of the endangered tiger beetle *Cicindela limbata albissima* (Coleoptera: Cicindelidae): evidence from mtDNA." Annals of the Entomological Society of America. Sept: 1108-1115.
- Morris, M. G. and W. E. Rispin (1988). "A beetle fauna of oolitic limestone grassland, and the responses of species to conservation management by different cutting regimes." Biological conservation **43**(2): 87-105.
- Mortimer, S. R., R. G. Booth, et al. (2002). "Effects of initial site management on the Coleoptera assemblages colonising newly established chalk grassland on ex-arable land." Biological Conservation **104**(3): 301-313.
- Morvan, N., Y. R. Delettre, et al. (1994). The distribution of Empididae (Diptera) in hedgerow network landscapes. Field margins: integrating agriculture and conservation, Coventry, Farnham.
- Mountford, E. P. (2002). "Fallen dead wood levels in the near-natural beech forest at La Tillaie reserve, Fontainebleau, France." Forestry **75**(2): 203-208.
- Mountford, E. P. and G. F. Peterken (2003). "Long-term change and implications for the management of wood-pastures: Experience over 40 years from Denny Wood, New Forest." Forestry **76**(1): 19-40.

- Mullen, K., O. Fahy, et al. (2003). "Ground flora and associated arthropod communities of forest road edges in Connemara, Ireland." Biodiversity and Conservation **12**(1): 87-101.
- Muller, M. M., M. Varama, et al. (2002). "Influence of insects on the diversity of fungi in decaying spruce wood in managed and natural forests." Forest Ecology and Management **166**(1-3): 165-181.
- Müller, M. M., M. Varama, et al. (2002). "Influence of insects on the diversity of fungi in decaying spruce wood in managed and natural forests." Forest Ecology and Management **166**(1-3): 165-181.
- Munks, S., K. Richards, et al. (2004). "Distribution, habitat and conservation of two threatened stag beetles, *Hoplogonus bornemisszai* and *H. vanderschoori* (Coleoptera: Lucanidae) in north-east Tasmania." Australian Zoologist **32**: 586-596.
- Murrell, D. J. and R. Law (2000). "Beetles in Fragmented Woodlands: A Formal Framework for Dynamics of Movement in Ecological Landscapes." Journal of Animal Ecology **69**(3): 471-483.
- Nappi, A., P. Drapeau, et al. (2004). "Salvage logging after wildfire in the boreal forest: Is it becoming a hot issue for wildlife?" Forestry Chronicle **80**(1): 67-74.
- Niemela, J., D. Langor, et al. (1993). "Effects of clear-cut harvesting on boreal ground-beetle assemblages (Coleoptera: Carabidae) in Western Canada." Conservation biology : the journal of the Society for Conservation Biology **7**(3): 551-561.
- Nilsen, H. C. and R. W. Larimore (1973). "Establishment of Invertebrate Communities on Log Substrates in the Kaskaskia River, Illinois." Ecology **54**(2): 366-374.
- Nilsson, S. G., M. Niklasson, et al. (2002). "Densities of large living and dead trees in old-growth temperate and boreal forests." Forest Ecology and Management **161**(1-3): 189-204.
- Nilsson, S. G., M. Niklasson, et al. (2003). "Erratum to "Densities of large living and dead trees in old-growth temperate and boreal forests"." Forest Ecology and Management **178**(3): 355-370.
- Nitterus, K., B. Gunnarsson, et al. (2004). "Insects reared from logging residue on clear-cuts." Entomologica Fennica **15**(1): 53-61.
- Nordén, B., T. Appelqvist, et al. (2002). "Diversity, ecology and conservation of wood-inhabiting ascomycetes in Sweden [Sporsäcksvampar i do?d ved - ma?ngfald, ekologi och naturva?rdsaspekter]." Svensk Botanisk Tidskrift **96**(3-4): 139-148.
- Norden, B., F. Gotmark, et al. (2004). "Dead wood in semi-natural temperate broadleaved woodland: contribution of coarse and fine dead wood, attached dead wood and stumps." Forest Ecology and Management **194**(1-3): 235-248.
- Nordén, B., M. Tönnberg, et al. (2004). "Dead wood in semi-natural temperate broadleaved woodland: Contribution of coarse and fine dead wood, attached dead wood and stumps." Forest Ecology and Management **194**(1): 235-248.
- Novotny, V., L. Cizek, et al. (2004). "Local Species Richness of Leaf-Chewing Insects Feeding on Woody Plants from One Hectare of a Lowland Rainforest." Conservation Biology **18**(1): 227-237.
- Nummelin, M. (1996). "The community structure of arthropods in virgin and managed sites in the Kibale Forest, Western Uganda." Tropical Ecology **37**(2): 203-213.

- Nyffeler, M. and K. D. Sunderland (2003). "Composition, abundance and pest control potential of spider communities in agroecosystems: a comparison of European and US studies." Agriculture, Ecosystems & Environment **95**(2-3): 579-612.
- Ødegaard, F. and F. Ødegaard (2004). "Species richness of phytophagous beetles in the tropical tree *Brosimum utile* (Moraceae): The effects of sampling strategy and the problem of tourists." Ecological Entomology **29**(1): 76-88.
- Okland, B. (1994). "Mycetophilidae (Diptera), an insect group vulnerable to forestry practices? A comparison of clearcut, managed and semi-natural spruce forests in southern Norway." Biodiversity and conservation **3**(1): 68-85.
- Okland, B. (1995). Diversity patterns of two insect groups within spruce forests of southern Norway, NORGES LANDBRUKSHOGSKOLE (NORWAY),.
- Okland, B. (1996). "Unlogged forests: important sites for preserving the diversity of mycetophilids (Diptera: Sciaroidea)." Biological Conservation **76**(3): 297-310.
- Økland, B. (1996). "A comparison of three methods of trapping saproxylic beetles." European Journal of Entomology **93**(2): 195-209.
- Okland, B., A. Bakke, et al. (1996). "What factors influence the diversity of saproxylic beetles? A multiscaled study from a spruce forest in southern Norway." Biodiversity and conservation **5**(1): 75-100.
- Økland, B., A. Bakke, et al. (1996). "What factors influence the diversity of saproxylic beetles? A multiscaled study from a spruce forest in southern Norway." Biodiversity and Conservation **5**(1): 75-100.
- Økland, B., F. Götmark, et al. (2005). "Regional diversity of mycetophilids (Diptera: Sciaroidea) in Scandinavian oak-dominated forests." Biological Conservation **121**(1): 9-20.
- Olberg, S., J. Andersen, et al. (2000). A survey of the saproxylic beetles (Coleoptera) in Dividalen, Troms county, northern Norway. Entomology, Melsomvik, Norway, Norwegian Entomological Society.
- Oliver, I., R. Mac Nally, et al. (2000). "Identifying performance indicators of the effects of forest management on ground-active arthropod biodiversity using hierarchical partitioning and partial canonical correspondence analysis." Forest Ecology and Management **139**(1-3): 21-40.
- O'Neal, M. E., E. L. Zontek, et al. (2005). "Ground predator abundance affects prey removal in highbush blueberry (*Vaccinium corymbosum*) fields and can be altered by aisle ground covers." Biocontrol **50**(2): 205-222.
- Orendt, C. (2000). "The chironomid communities of woodland springs and spring brooks, severely endangered and impacted ecosystems in a lowland region of eastern Germany (Diptera: Chironomidae)." Journal of Insect Conservation **4**(2): 79-91.
- Oslash and d. F. Species richness of phytophagous beetles in the tropical tree *Brosimum utile* (Moraceae): the effects of sampling strategy and the problem of tourists. Ecological Entomology. **29**: 76-88.
- Ovcharov, D. and D. Doychev (2001). Insect pests of the order Coleoptera in recently created pine plantations on non-typical habitats. Balkan scientific conference; Study, conservation and utilisation of forest resources, Sofia, Bulgaria, Academic Publishers.

Ozanne, C. M. P., M. R. Speight, et al. (2000). "Isolated trees and forest patches: Patterns in canopy arthropod abundance and diversity in *Pinus sylvestris* (Scots Pine)." Forest Ecology and Management **137**(1-3): 53-63.

Painter, D. (1999). "Macroinvertebrate distributions and the conservation value of aquatic Coleoptera, Mollusca and Odonata in the ditches of traditionally managed and grazing fen at Wicken Fen, UK." Journal of applied ecology **36**(1): 33-36.

Palmer, S. C. F., R. J. Mitchell, et al. (2004). "Regeneration failure in Atlantic oakwoods: The roles of ungulate grazing and invertebrates." Forest Ecology and Management **192**(2-3): 251-265.

Parsons, L. S. and W. H. Lear (2001). "Climate variability and marine ecosystem impacts: a North Atlantic perspective." Progress In Oceanography **49**(1-4): 167-188.

Parsons, T. (1995). "The saproxylic invertebrates of Bincombe Hill, Crewkerne, Somerset." Somerset fauna & flora **1**.

Patterson, I. J., J. G. Ollason, et al. (1995). "Bird populations in upland spruce plantations in northern Britain." Forest Ecology and Management **79**(1-2): 107-131.

Pavett, B. L. a. P. M. (2000). "Dinefwr Park saproxylic invertebrates." CCW contract science report **353**.

Paviour-Smith, K. and C. A. Elbourn (1992). A quantitative study of the fauna of small dead and dying wood in living trees in Wytham Woods, near Oxford. Dead wood matters: the ecology and conservation of saproxylic invertebrates in Britain, Peterborough.

Pavuk, D. M., F. F. Purrington, et al. (1997). "Ground beetle (Coleoptera: Carabidae) activity density and community composition in vegetationally diverse corn agroecosystems." American midland naturalist **138**(1): 14-28.

Pechacek, P., A. Kristin, et al. (2004). "Comparative diets of adult and young three-toed woodpeckers in a European alpine forest community." Journal of Wildlife Management **68**(3): 683-693.

Pedersen, B. S. and B. McCune (2002). "A non-invasive method for reconstructing the relative mortality rates of trees in mixed-age, mixed-species forests." Forest Ecology and Management **155**(1-3): 303-314.

Peeters, P. J., J. Read, et al. (2001). "Variation in the guild composition of herbivorous insect assemblages among co-occurring plant species." Austral Ecology **26**(4): 385-399.

Peltonen, M. (1999). "Windthrows and dead-standing trees as bark beetle breeding material at forest-clearcut edge." Scandinavian Journal of Forest Research **14**(6): 505-511.

Peltonen, M. and K. Heliövaara (1998). "Incidence of *Xylechinus pilosus* and *Cryphalus saltuarius* (Scolytidae) in forest-clearcut edges." Forest Ecology and Management **103**(2-3): 141-147.

Peltonen, M., K. Heliövaara, et al. (1997). "Forest insects and environmental variation in stand edges." Silva Fennica **31**(2): 129-141.

Perfecto, I., J. Vandermeer, et al. (1997). "Arthropod biodiversity loss and the transformation of a tropical agro-ecosystem." Biodiversity and conservation **6**(7): 935-945.

Perison, D., J. Phelps, et al. (1997). "The effects of timber harvest in a South Carolina blackwater bottomland." Forest Ecology and Management **90**(2-3): 171-185.

- Perkins, A. J., M. J. Whittingham, et al. (2000). "Habitat characteristics affecting use of lowland agricultural grassland by birds in winter." Biological Conservation **95**(3): 279-294.
- Peterken, G. F. (1999). "Applying natural forestry concepts in an intensively managed landscape." Global Ecology and Biogeography **8**(5): 321-328.
- Peterken, G. F. (2001). "Ecological effects of introduced tree species in Britain." Forest Ecology and Management **141**(1-2): 31-42.
- Peterken, G. F. and J. L. Francis (1999). "Open spaces as habitats for vascular ground flora species in the woods of central Lincolnshire, UK." Biological Conservation **91**(1): 55-72.
- Petit, S., K. Haysom, et al. (2003). "Habitat-based models for predicting the occurrence of ground- beetles in arable landscapes: two alternative approaches." Agriculture Ecosystems & Environment **95**(1): 19-28.
- Petranka, J. W., M. P. Brannon, et al. (1994). "Effects of timber harvesting on low elevation populations of southern Appalachian salamanders." Forest Ecology and Management **67**(1-3): 135-147.
- Pettersson, R. B., J. P. Ball, et al. (1995). "Invertebrate communities in boreal forest canopies as influenced by forestry and lichens with implications for passerine birds." Biological Conservation **74**(1): 57-63.
- Pfarr, U. and J. Schrammel (1991). "Dead Woody Material of Norway Spruce Cause for Strained Relations between Nature Conservation and Forestry." Forstwissenschaftliches Centralblatt **110**(2): 128-134.
- Piper, R. W., S. G. Compton, et al. (2001). "The species status of *Cathormiocerus britannicus*, an endemic, endangered British Weevil." Biological Conservation **101**(1): 9-13.
- Pitkanen, S. (1998). "The use of diversity indices to assess the diversity of vegetation in managed boreal forests." Forest Ecology and Management **112**(1-2): 121-137.
- Pollard, E. M. L. H. J. N. G.-D. (1987). "Management guidelines for the conservation of invertebrates, especially butterflies, in plantation woodland." Huntingdon : Institute of Terrestrial Ecology.
- Poole, A., M. Gormally, et al. (2003). "The flora and carabid beetle fauna of a mature and regenerating semi-natural oak woodland in south-east Ireland." Forest Ecology and Management **177**(1-3): 207-220.
- Potting, R. P. J., J. N. Perry, et al. (2005). "Insect behavioural ecology and other factors affecting the control efficacy of agro-ecosystem diversification strategies." Ecological Modelling **182**(2): 199-216.
- Progar, R. A. and T. D. Schowalter (2002). "Canopy arthropod assemblages along a precipitation and latitudinal gradient among Douglas-fir *Pseudotsuga menziesii* forests in the Pacific Northwest of the United States." Ecography **25**(2): 129-138.
- Progar, R. A., T. D. Schowalter, et al. (2000). "Respiration from coarse woody debris as affected by moisture and saprotroph functional diversity in Western Oregon." Oecologia **124**(3): 426-431.
- Punttila, P. and Y. Haila (1996). "Colonisation of a burned forest by ants in the Southern Finnish boreal forest." Silva Fennica **30**(4): 421-435.
- Quinn, J. M., I. K. G. Boothroyd, et al. (2004). "Riparian buffers mitigate effects of pine plantation logging on New Zealand streams: 2. Invertebrate communities." Forest Ecology and Management **191**(1-3): 129-146.

Radford, J. Q., A. F. Bennett, et al. (2005). "Landscape-level thresholds of habitat cover for woodland-dependent birds." Biological Conservation **124**(3): 317-337.

Ranius, T. (2002). "Influence of stand size and quality of tree hollows on saproxylic beetles in Sweden." Biological Conservation **103**(1): 85-91.

Ranius, T. (2002). "Osmoderma eremita as an indicator of species richness of beetles in tree hollows." Biodiversity and conservation **11**(5): 931-941.

Ranius, T. (2002). "Population ecology and conservation of beetles and pseudoscorpions living in hollow oaks in Sweden." Animal Biodiversity and Conservation **25**(1): 53-68.

Ranius, T. and N. Jansson (2000). "The influence of forest regrowth, original canopy cover and tree size on saproxylic beetles associated with old oaks." Biological Conservation **95**(1): 85-94.

Ranius, T. and N. Jansson (2002). "A comparison of three methods to survey saproxylic beetles in hollow oaks." Biodiversity and conservation **11**(10): 1759-1771.

Ranius, T., B. G. Jonsson, et al. (2004). "Modeling dead wood in Fennoscandian old-growth forests dominated by Norway spruce." Canadian Journal of Forest Research **34**(5): 1025-1034.

Ranius, T. and O. Kindvall (2004). "Modelling the amount of coarse woody debris produced by the new biodiversity-oriented silvicultural practices in Sweden." Biological Conservation **119**(1): 51-59.

Ranius, T., O. Kindvall, et al. (2003). "Modelling dead wood in Norway spruce stands subject to different management regimes." Forest Ecology and Management **182**(1-3): 13-29.

Ranius, T. and S. G. Nilsson (1997). "Habitat of *Osmoderma eremita* Scop. (Coleoptera: Scarabaeidae), a beetle living in hollow trees." Journal of Insect Conservation **1**(4): 193-204.

Ratcliffe, P. R. and G. F. Peterken (1995). "The potential for biodiversity in British upland spruce forests." Forest Ecology and Management **79**(1-2): 153-160.

Rauh, J. and M. Schmitt (1991). "Methods and Results of Research on Dead Woody Material on Natural-Forest Reserves." Forstwissenschaftliches Centralblatt **110**(2): 114-127.

Reemer, M. (2005). "Saproxylic hoverflies benefit by modern forest management (Diptera: Syrphidae)." Journal of Insect Conservation **9**(1): 49-59.

Ribera, I. Biogeography and conservation of Iberian water beetles - graphical techniques, smoothing and searching for what makes relationships work. Biological Conservation. **92**: 131-150.

Richard, E., F. Gosselin, et al. (2003). Short-term and Mid-term Response of Ground Beetle Communities (Coleoptera, Carabidae) to Disturbance by Regeneration Felling. History and forest biodiversity: challenges for conservation; Forest biodiversity lessons from history for conservation, Leuven, Belgium, Wallingford.

Romero-Alcaraz, E. and J. M. Avila (2000). "Landscape heterogeneity in relation to variations in epigeic beetle diversity of a Mediterranean ecosystem. Implications for conservation." Biodiversity and conservation **9**(7): 985-1005.

- Roth, D. S., I. Perfecto, et al. (1994). "The Effects of Management Systems on Ground-Foraging Ant Diversity in Costa Rica." Ecological Applications **4**(3): 423-436.
- Rotheray, G. E., G. Hancock, et al. (2001). "The biodiversity and conservation of saproxylic Diptera in Scotland." Journal of Insect Conservation **5**(2): 77-85.
- Rotheray, G. E. and I. MacGowan (2000). "Status and breeding sites of three presumed endangered Scottish saproxylic syrphids (Diptera, Syrphidae)." Journal of Insect Conservation **4**(4): 215-223.
- Rozas, V. (2004). "A dendroecological reconstruction of age structure and past management in an old-growth pollarded parkland in northern Spain." Forest Ecology and Management **195**(1-2): 205-219.
- Rukke, B. A. (2000). "Effects of habitat fragmentation: increased isolation and reduced habitat size reduces the incidence of dead wood fungi beetles in a fragmented forest landscape." Ecography **23**(4): 492-502.
- Rukke, B. A. and F. Midtgaard (1998). "The importance of scale and spatial variables for the fungivorous beetle *Bolitophagus reticulatus* (Coleoptera, Tenebrionidae) in a fragmented forest landscape." Ecography **21**(6): 561-572.
- Saint-Germain, M., P. Drapeau, et al. (2004). "Comparison of Coleoptera assemblages from a recently burned and unburned black spruce forests of northeastern North America." Biological Conservation **118**(5): 583-592.
- Saint-Germain, M., M. Larrivee, et al. "Short-term response of ground beetles (Coleoptera: Carabidae) to fire and logging in a spruce-dominated boreal landscape." Forest Ecology and Management **In Press, Corrected Proof**.
- Schiegg, K. (2000). "Are there saproxylic beetle species characteristic of high dead wood connectivity?" Ecography **23**(5): 579-587.
- Schiegg, K. (2000). "Effects of dead wood volume and connectivity on saproxylic insect species diversity." Écoscience (Sainte-Foy, Quebec) **Écoscience** **7**(3): 290-298.
- Schiegg, K. (2001). "Saproxylic insect diversity of beech: limbs are richer than trunks." Forest ecology and management **149**(1/3): 295-304.
- Schiegg, K. and K. Schiegg (2003). "Erratum: Saproxylic insect diversity of beech: Limbs are richer than trunks (Forest Ecology and Management (2001) 149:1-3 (295-304)PII: S037811270000 5636)." Forest Ecology and Management **175**(1-3): 589.
- Schlaghamersky, J. (1999). The impact of flooding on saproxylic beetles and ants in South Moravian floodplain forests. Central European workshop on soil zoology; Soil zoology in Central Europe, Ceske Budejovice, Czech Republic, Ceske Budejovice.
- Schlichter, D. (1975). "Produktion oder ubernahme von schutzstoffen als ursache des Nesselschutzes von Anemonenfischen?" Journal of Experimental Marine Biology and Ecology **20**(1): 49-61.
- Schmidl, J. and H. Bussler (2004). "Ecological Guilds of Xylobiontic (Saproxylic) Beetles in Germany and Their Use in Landscape Ecological Surveys." Naturschutz Und Landschaftsplanung. **36**(7): 202-218.
- Schops, K., S. D. Wratten, et al. (1999). "Life cycle, behaviour and conservation of the large endemic weevil, *Hadramphus spinipennis* on the Chatham Islands, New Zealand." New Zealand Journal of Zoology **26**(1): 55-66.

- Schroeder, L. M., J. Weslien, et al. (1999). "Attacks by bark- and wood-boring Coleoptera on mechanically created high stumps of Norway spruce in the two years following cutting." Forest Ecology and Management **123**(1): 21-30.
- Schröter, H. (2003). "Biodiversity and integrated forest pest control [Biodiversität und integrierter Waldschutz]." Gesunde Pflanzen **55**(4): 93-97.
- Seastedt, T. R. and D. A. Crossley, Jr. (1981). "Microarthropod Response Following Cable Logging and Clear-Cutting in the Southern Appalachians." Ecology **62**(1): 126-135.
- Sekercioglu, C. H., P. R. Ehrlich, et al. (2002). "Disappearance of Insectivorous Birds from Tropical Forest Fragments." Proceedings of the National Academy of Sciences of the United States of America **99**(1): 263-267.
- Shackleton, C. M. (1998). "Annual production of harvestable deadwood in semi-arid savannas, South Africa." Forest Ecology and Management **112**(1-2): 139-144.
- Shorrocks, B. (1975). "The Distribution and Abundance of Woodland Species of British Drosophila (Diptera: Drosophilidae)." The Journal of Animal Ecology **44**(3): 851-864.
- Siira-Pietikainen, A., J. Haimi, et al. (2001). "Responses of decomposer community to root-isolation and addition of slash." Soil Biology and Biochemistry **33**(14): 1993-2004.
- Siira-Pietikainen, A., J. Haimi, et al. (2003). "Short-term responses of soil macroarthropod community to clear felling and alternative forest regeneration methods." Forest Ecology and Management **172**(2-3): 339-353.
- Siitonen, J. (1994). "Decaying wood and saproxylic Coleoptera in two old spruce forests: a comparison based on two sampling methods." Annales Zoologici Fennici **31**(1): 89-95.
- Siitonen, J. (1999). Forest management, coarse woody debris and saproxylic organisms: Fennoscandian boreal forests as an example. Ecology of coarse woody debris in boreal forests, Oxford.
- Siitonen, J. and S. L. Habitat requirements and conservation of *Pytho kolwensis*, a beetle species of old-growth boreal forest. Biological Conservation **94**: 211-220.
- Siitonen, J., P. Martikainen, et al. (1996). "New faunistic records of threatened saproxylic Coleoptera, Diptera, Heteroptera, Homoptera and Lepidoptera from the Republic of Karelia, Russia." Entomologica Fennica **7**(2): 69-76.
- Siitonen, J., P. Martikainen, et al. (2000). "Coarse woody debris and stand characteristics in mature managed and old-growth boreal mesic forests in southern Finland." Forest Ecology and Management **128**(3): 211-225.
- Siitonen, J., R. Penttilä, et al. (1999). Coarse woody debris, polyporous fungi and saproxylic insects in an old-growth spruce forest in Vodlozero National Park, Russian Karelia. Ecology of coarse woody debris in boreal forests, Oxford.
- Siitonen, J. and L. Saaristo (2000). "Habitat requirements and conservation of *Pytho kolwensis*, a beetle species of old-growth boreal forest." Biological Conservation **94**(2): 211-220.
- Sikes, D. S. and C. J. Raithel (2002). "A review of hypotheses of decline of the endangered American burying beetle (Silphidae: *Nicrophorus americanus* Olivier)." Journal of Insect Conservation **6**(2): 103-113.

- Simandl, J. (1993). "Canopy arthropods on Scots pine: influence of season and stand age on community structure and the position of sawflies (Diprionidae) in the community." Forest Ecology and Management **62**(1-4): 85-98.
- Similä, M., J. Kouki, et al. (2003). "Saproxylic beetles in managed and seminatural Scots pine forests: quality of dead wood matters." Forest Ecology and Management **174**(1-3): 365-381.
- Similä, M., J. Kouki, et al. (2002). "Conservation of beetles in boreal pine forests: the effects of forest age and naturalness on species assemblages." Biological Conservation **106**(1): 19-27.
- Similä, M., J. Kouki, et al. (2002). "Conservation of beetles in boreal pine forests: The effects of forest age and naturalness on species assemblages." Biological Conservation **106**(1): 19-27.
- Sippola, A.-L. (2001). Forest structure and biodiversity in northern boreal forests: Effects of regeneration cutting on flying beetles and wood-decomposing fungi, HELSINGIN YLIOPISTO (FINLAND),.
- Sippola, A.-L., R. Kallio, et al. (1998). "Amount and quality of coarse woody debris in natural and managed coniferous forests near the timberline in Finnish Lapland." Scandinavian Journal of Forest Research **13**(2): 204-214.
- Sippola, A.-L. and P. Renvall (1999). "Wood-decomposing fungi and seed-tree cutting: A 40-year perspective." Forest Ecology and Management **115**(2-3): 183-201.
- Sippola, A.-L., J. Siitonen, et al. (2002). "Beetle diversity in timberline forests: A comparison between old-growth and regeneration areas in Finnish Lapland." Annales Zoologici Fennici **39**(1): 69-86.
- Skidmore, P. (2003). "Saproxylic insect survey of the Virginia Water and Bishopsgate areas of Windsor Park, 2002-2003." English Nature research reports **514**.
- Small, E. C., J. P. Sadler, et al. (2002). "Carabid beetle assemblages on urban derelict sites in Birmingham, UK." Journal of Insect Conservation **6**(4): 233-246.
- Smallidge, P. J. and D. J. Leopold (1997). "Vegetation management for the maintenance and conservation of butterfly habitats in temperate human-dominated landscapes." Landscape and Urban Planning **38**(3-4): 259-280.
- Smart, R., M. J. Whiting, et al. (2005). "Lizards and landscapes: integrating field surveys and interviews to assess the impact of human disturbance on lizard assemblages and selected reptiles in a savanna in South Africa." Biological Conservation **122**(1): 23-31.
- Smith, K. W. (1997). "Nest site selection of the great spotted woodpecker *Dendrocopos major* in two oak woods in Southern England and its implications for woodland management." Biological Conservation **80**(3): 283-288.
- Smith, V. G. (1928). "Animal Communities of a Deciduous Forest Succession." Ecology **9**(4): 479-500.
- Species, P. s. T. f. E. "Proceedings of the second pan-European conference on Saproxylic Beetles."
- Speight, M. C. D. (1989). "Saproxylic invertebrates and their conservation." Nature and environment (Council of Europe) **42**.
- Standish, R. J. (2004). "Impact of an invasive clonal herb on epigeic invertebrates in forest remnants in New Zealand." Biological Conservation **116**(1): 49-58.

- Stephen, F. M. and L. E. Browne (2000). "Application of Eliminate parasitoid food to boles and crowns of pines (Pinaceae) infested with *Dendroctonus frontalis* (Coleoptera: Scolytidae)." Canadian entomologist **132**(6): 983-985.
- Stokland, J. N. (1997). "Representativeness and efficiency of bird and insect conservation in Norwegian boreal forest reserves [Representatividad y eficiencia en al conservacion de aves e insectos en la reservas de bosque boreal de Noruega]." Conservation Biology **11**(1): 101-111.
- Stoll, P., J. Weiner, et al. (1994). "Growth Variation in a Naturally Established Population of *Pinus Sylvestris*." Ecology **75**(3): 660-670.
- Stouffer, P. C. and R. O. Bierregaard, Jr. (1995). "Use of Amazonian Forest Fragments by Understory Insectivorous Birds." Ecology **76**(8): 2429-2445.
- Strehlow, K., J. S. Bradley, et al. (2002). "Short term impacts of logging on invertebrate communities in jarrah forests in south-west Western Australia." Forest Ecology and Management **162**(2-3): 165-184.
- Stuart-Smith, A. K. and J. P. Hayes (2003). "Influence of residual tree density on predation of artificial and natural songbird nests." Forest Ecology and Management **183**(1-3): 159-176.
- Stubbs, A. (1999). "Beavers and saproxylic flies [1]." British Wildlife **10**(3): 219.
- Summers, R. W., R. A. Mavor, et al. (1999). "The structure of ancient native pinewoods and other woodlands in the Highlands of Scotland." Forest Ecology and Management **119**(1-3): 231-245.
- Sverdrup-Thygeson, A. (2001). "Can 'continuity indicator species' predict species richness or red-listed species of saproxylic beetles?" Biodiversity and Conservation **10**(5): 815-832.
- Sverdrup-Thygeson, A. and R. A. Ims (2002). "The effect of forest clearcutting in Norway on the community of saproxylic beetles on aspen." Biological Conservation **106**(3): 347-357.
- Sverdrup-Thygeson, A. and R. A. Ims (2002). "Erratum: The effect of forest clearcutting in Norway on the community of saproxylic beetles on aspen (Biological Conservation (2000) 108:3(377-378) PII: S0006320701002610)." Biological Conservation **108**(3): 377-378.
- Taboada, A., D. J. Kotze, et al. (2004). "Carabid beetle occurrence at the edges of oak and beech forests in NW Spain." European Journal of Entomology **101**(4): 555-563.
- Thomas, F., J.-F. Guegan, et al. (2000). "Parasites and host life-history traits: implications for community ecology and species co-existence." International Journal for Parasitology **30**(5): 669-674.
- Thomas, J. A. and M. G. Morris (1994). "Patterns, Mechanisms and Rates of Extinction among Invertebrates in the United Kingdom." Philosophical Transactions: Biological Sciences **344**(1307, Estimating Extinction Rates: Sir Joseph Banks Anniversary Meeting): 47-54.
- Thomas, R. C., K. J. Kirby, et al. (1997). "The conservation of a fragmented ecosystem within a cultural landscape--The case of ancient woodland in England." Biological Conservation **82**(3): 243-252.
- Thomas, S. R., R. Noordhuis, et al. (2002). "Botanical diversity of beetle banks: Effects of age and comparison with conventional arable field margins in southern UK." Agriculture, Ecosystems & Environment **93**(1-3): 403-412.

- Thunes, K. H., I. Gjerde, et al. (2000). "Diversity of coleoptera of the bracket fungus *Fomitopsis pinicola* in a Norwegian spruce forest." *Biodiversity and Conservation* **9**(6): 833-852.
- Tidon, R., D. F. Leite, et al. (2003). "Impact of the colonisation of *Zaprionus* (Diptera, Drosophilidae) in different ecosystems of the Neotropical Region: 2 years after the invasion." *Biological Conservation* **112**(3): 299-305.
- Timoney, K. P. and G. Peterson (1996). "Failure of natural regeneration after clearcut logging in Wood Buffalo National Park, Canada." *Forest Ecology and Management* **87**(1-3): 89-105.
- Timothy, L. K. (1991). The impact of coniferous afforestation on surface water quality in the Kielder Area, Northumberland., Newcastle upon Tyne.
- Truscott, A. M., R. J. Mitchell, et al. (2004). "The expansion of native oakwoods into conifer cleared areas through planting." *Forest Ecology and Management* **193**(3): 335-343.
- Turin, H. and P. J. d. Boer (1988). "Changes in the distribution of carabid beetles in the Netherlands since 1880. II. Isolation of habitats and long-term time trends in the occurrence of carabid species with different powers of dispersal (Coleoptera, Carabidae)." *Biological conservation* **44**(3): 179-200.
- Uliczka, H. (2003). "Nature conservation efforts by forest owners - Intentions and practice in a Swedish case study." *Silva Fennica* **37**(4): 459-475.
- Ulyshen, M. D., J. L. Hanula, et al. (2004). "Spatial and temporal patterns of beetles associated with coarse woody debris in managed bottomland hardwood forests." *Forest Ecology and Management* **199**(2-3): 259-272.
- Vaisanen, R., O. Bistrom, et al. (1993). "Sub-cortical Coleoptera in dead pines and spruces: is primeval species composition maintained in managed forests?" *Biodiversity and conservation* **2**(2): 95-113.
- Valladares, L. F., J. Garrido, et al. (2002). "The assemblages of aquatic Coleoptera from shallow lakes in the northern Iberian Meseta: Influence of environmental variables." *European Journal of Entomology* **99**(3): 289-298.
- Vasiliauskas, R., A. Vasiliauskas, et al. (2004). "Dead trees and protected polypores in unmanaged north-temperate forest stands of Lithuania." *Forest Ecology and Management* **193**(3): 355-370.
- Virkkala, R. (2004). "Bird species dynamics in a managed southern boreal forest in Finland." *Forest Ecology and Management* **195**(1-2): 151-163.
- Wallace, H. R. (1953). "The Ecology of the Insect Fauna of Pine Stumps." *The Journal of Animal Ecology* **22**(1): 154-171.
- Wardell-Johnson, G. and P. Horwitz (1996). "Conserving biodiversity and the recognition of heterogeneity in ancient landscapes: a case study from southwestern Australia." *Forest Ecology and Management* **85**(1-3): 219-238.
- Warren, G. L. (1956). "The Effect of Some Site Factors on the Abundance of *Hypomolyx Piceus* (Coleoptera: Curculionidae)." *Ecology* **37**(1): 132-139.
- Warren, M. S., E. Pollard, et al. (1986). "Annual and Long-Term Changes in a Population of the Wood White Butterfly *Leptidea sinapis*." *The Journal of Animal Ecology* **55**(2): 707-719.
- Warriner, M. D. (2004). "Cerambycidae. Occurrence and Conservation Status of the Milkweed Beetle *Tetraopes texanus* Horn (Coleoptera: Cerambycidae) in Arkansas." *Coleopterists Bulletin* **58**: 567-568.
- Warriner, M. D. (2004). "Occurrence and conservation status of the milkweed beetle *Tetraopes texanus* Horn (Coleoptera: Cerambycidae) in Arkansas." *Coleopterists bulletin* **58**: 567-568.

- Watson, J. E. M., R. J. Whittaker, et al. (2004). "Habitat structure and proximity to forest edge affect the abundance and distribution of forest-dependent birds in tropical coastal forests of southeastern Madagascar." Biological Conservation **120**(3): 311-327.
- Watt, A. S. (1923). "On the Ecology of British Beechwoods with Special Reference to Their Regeneration." The Journal of Ecology **11**(1): 1-48.
- Weikel, J. M. and J. P. Hayes (1999). "The foraging ecology of cavity-nesting birds in young forests of the Northern Coast Range of Oregon." Condor **101**(1): 58-66.
- Welsh, R. C. (1995). Native and non-native trees as habitats for phytophagous and saproxylic invertebrates. Native and non-native in British forestry, Coventry, Institute of Chartered Foresters.
- Wermelinger, B. (2004). "Ecology and management of the spruce bark beetle *Ips typographus* - A review of recent research." Forest Ecology and Management **202**(1-3): 67-82.
- Wermelinger, B., P. Duelli, et al. (2002). "Dynamics of saproxylic beetles (Coleoptera) in windthrow areas in alpine spruce forests." Forest Snow and Landscape Research **77**(1-2): 133-148.
- Werner, S. M. and K. F. Raffa (2000). "Effects of forest management practices on the diversity of ground-occurring beetles in mixed northern hardwood forests of the Great Lakes Region." Forest Ecology and Management **139**(1-3): 135-155.
- Weslien, J. and L. Martin Schroeder (1999). "Population levels of bark beetles and associated insects in managed and unmanaged spruce stands." Forest Ecology and Management **115**(2-3): 267-275.
- Whitehead, P. F. (1997). "Beetle faunas of the European angiosperm Urwald: Problems and complexities." Biologia - Section Zoology **52**(2): 147-152.
- Whitehead, P. F. (2005). "The saproxylic quality Coleoptera of Brockworth Great Park, Gloucestershire, the issue of site ranking in Britain and new concepts of Arboreal Niche Diversity and Index Reconciliation." Entomologist's Gazette **56**(1): 35-46.
- Wiedenmann, R. N., R. R. Diaz, et al. (2004). "Ground beetle (Coleoptera: Carabidae) assemblages in mulched and non-mulched garden plots." Journal of the Kansas Entomological Society **77**: 99-109.
- Wikars, L.-O. and J. Schimmel (2001). "Immediate effects of fire-severity on soil invertebrates in cut and uncut pine forests." Forest Ecology and Management **141**(3): 189-200.
- Williams, A. B. (1936). "The Composition and Dynamics of a Beech-Maple Climax Community." Ecological Monographs **6**(3): 317-408.
- Williams, M. R., I. Abbott, et al. (2001). "Recovery of bird populations after clearfelling of tall open eucalypt forest in Western Australia." Journal of Applied Ecology **38**(5): 910-920.
- Willott, S. J. (1999). "The Effects of Selective Logging on the Distribution of Moths in a Bornean Rainforest." Philosophical Transactions: Biological Sciences **354**(1391, Changes and Disturbance in Tropical Rainforest in South-East Asia): 1783-1790.
- Winter, T. (1992). Deadwood - is it a threat to commercial forestry? Dead wood matters: the ecology and conservation of saproxylic invertebrates in Britain, Peterborough.

- Wolters, V. (1991). "Biological processes in two beech forest soils treated with simulated acid rain--a laboratory experiment with *Isotoma tigrina* (insecta, collembola)." Soil Biology and Biochemistry **23**(4): 381-390.
- Woodcock, B. A., A. D. Watt, et al. (2003). "Influence of management type on Diptera communities of coniferous plantations and deciduous woodlands." Agriculture Ecosystems & Environment **95**(2-3): 443-452.
- Work, T. T., J. R. Spence, et al. (2004). "Stand composition and structure of the boreal mixedwood and epigeaic arthropods of the Ecosystem Management Emulating Natural Disturbance (EMEND) landbase in northwestern Alberta." Canadian Journal of Forest Research **34**(2): 417-430.
- Xuluc-Tolosa, F. J., H. F. M. Vester, et al. (2003). "Leaf litter decomposition of tree species in three successional phases of tropical dry secondary forest in Campeche, Mexico." Forest Ecology and Management **174**(1-3): 401-412.
- Zaragoza-Caballero, S., F. A. Noguera, et al. (2003). "Diversity of Lycidae, Phengodidae, Lampyridae, and Cantharidae (Coleoptera) in a tropical dry forest region in Mexico: Sierra de Huautla, Morelos." Pan-Pacific entomologist **79**(1): 23-37.
- Zavala, M. A. and J. A. Oria (1995). "Preserving biological diversity in managed forests: a meeting point for ecology and forestry." Landscape and Urban Planning **31**(1-3): 363-378.
- Zilihona, I. J. E. and N. M. Coleopteran diversity and abundance in different habitats near Kihansi waterfall, in the Udzungwa Mountains, Tanzania. Biodiversity and Conservation. **10**: 769-777.
- Zimmer, M. (2002). "Is decomposition of woodland leaf litter influenced by its species richness?" Soil Biology and Biochemistry **34**(2): 277-284.